

Museum Victoria Science Report

Number 11, 2008

<https://doi.org/10.24199/j.mvsr.2008.11>



Decapod Crustacea of the continental margin
of southwestern and central Western Australia:
preliminary identifications of 524 species from
FRV *Southern Surveyor* voyage SS10-2005

Gary C. B. Poore, Anna W. McCallum and Joanne Taylor

Decapod Crustacea of the continental margin of southwestern and central Western Australia: preliminary identifications of 524 species from FRV *Southern Surveyor* voyage SS10-2005

GARY C. B. POORE, ANNA W. MCCALLUM AND JOANNE TAYLOR

Museum Victoria, GPO Box 666E, Melbourne, Victoria 3001, Australia
(gpoore@museum.vic.gov.au)

Abstract

Poore, G.C.B., McCallum, A.S., and Taylor, J. 2008. Decapod Crustacea of the continental margin of southwestern and central Western Australia: preliminary identifications of 524 species from FRV *Southern Surveyor* voyage SS10-2005. *Museum Victoria Science Reports* 11: 1–106.
A collection of Dendrobranchiata (44 species), Achelata (4 species), Anomura (127 species), Astacidea (4 species), Brachyura (227 species), Caridea (88 species), Polychelida (5 species), Stenopodidea (2 species) and Thalassinidea (23 species) from shelf edge and slope depths of south-western Australia is reported. Seventy-seven families are represented. Thirty-three per cent (175) of all species are suspected to be new species, eight per cent are new records for Australia, and a further 25% newly recorded for southern Western Australia.

Contents

Introduction.....	1
Methods	2
Presentation of taxonomic results	3
Taxonomic results and commentary	3
Invitation.....	4
Dendrobranchiata.....	12
Achelata	16
Anomura	17
Astacidea.....	34
Brachyura.....	35
Caridea.....	78
Polychelida.....	91
Stenopodidea.....	92
Thalassinidea	92
Acknowledgements.....	97
References.....	97
Index to families and genera	104

Introduction

The offshore fauna of southwestern Australia is poorly known relative to that of some other regions of Australia. Population centres in southeastern Australia and offshore oil and gas resources in the northwest have in different ways provided foci for exploration and some biological characterisation along the continental margins. Marine exploration in the southwest on the other hand has been confined to coastal and shallow-water environments, particularly in a series of taxonomic workshops in Albany, Rottnest Island and Esperance. Decapod crustaceans from these regions were reviewed by Morgan and Jones (1991) and Jones and Morgan (1993). Crustacean collections made offshore and now residing in the Western Australian Museum are not extensive, those from the cruises of the FV *Davena* (1960) and HMAS *Diamantina* (1960s) being the most significant.

All of this is ironic because the first ever illustrations by Europeans of Australian marine animals were published in 1703 by the privateer William Dampier (1651–1715) (Dampier, 1703). Many of the shore collections made by François Peron and colleagues during the 1802 visit of the *Naturaliste* and *Geographe* to Australia were made in southwestern Australia (Milne Edwards, 1837). Later foreign expeditions also targeted the southwest (Balss, 1935).

During compilation of records for a guide to identification of southern Australian decapod Crustacea (Poore, 2004) it emerged that the southwest was less well known than the southeast of Australia. This impression is borne out by an analysis of the distribution of species along the southern Australian coast (O'Hara and Poore, 2000). These authors discovered that species composition varied with both latitude and longitude. Species richness was relatively constant from east to west but graded with latitude from high in the warm temperate regions around Perth and Sydney to low in cool-temperate southern Tasmania. They concluded that history as well as ecological hypotheses explain the latitudinal gradient of marine species richness in southern Australia, not the least being the invasion of the southwestern margin by animals of Indo-West Pacific origin.

Bioregionalisation of southwestern Australia depends now on geophysical surrogates and patterns in the distribution of fishes of the shelf and continental slope (Last et al., 2005). Three bioregions have been recognised along the WA coast with two intermediate regions in between: the Northwest Province, Central Western Transition Zone, Central Western Province, Southwestern Transition Zone and Southern Province.

The results presented here are part of a project mounted largely by CSIRO Marine and Atmospheric Research (CMAR) and Museum Victoria entitled “Mapping benthic ecosystems on the deep continental shelf and slope in Australia’s South West Region” to understand evolution and biogeography and support implementation of the SW Regional Marine Plan and

Commonwealth Marine Protected Areas". The field work addresses four primary objectives:

"1) test hypotheses on the evolution and biogeography of Australia's biodiversity, in particular relating to species composition, distribution patterns and taxonomic surrogacy

2) validate and refine CSIRO's optimised methodology for mapping deep water benthic ecosystems on the western continental margin and in sub-tropical locations to enhance its application to natural resource management at a national scale

3) document the benthic biodiversity and identify areas of high conservation values in the context of Commonwealth MPA declaration

4) validate, and permit refinement of, a marine bioregionalisation during the development of the SW Regional Marine Plan by the National Oceans Office."

This report deals only with the crustacean Order Decapoda, one of the taxa chosen to test these hypothesis. It first outlines briefly where and how the new material was collected. Next, data on taxonomy and distribution associated with each taxon identified are presented with brief comments. The purpose of publishing summarised information is to alert taxonomists to this essentially new and previously undescribed fauna and to provide access to data for a distributional analysis of the region.




Figure 1. The survey area showing positions of sampling sites. At 11 sites between Albany and Exmouth samples were taken at depths of 100 and 400 m (black stars), and at 7 transects on special features at intended depths of 100, 200, 400, 700 and 1000 m (red rectangles).

Methods

Sampling program. The data for this project were collected during two surveys undertaken from FRV *Southern Surveyor*, a 67 m converted stern-trawler. The first was completed in July–August 2005 when all the survey sites were mapped using multibeam acoustics, surveyed with a towed, high-resolution video system and sampled with sediment grabs. The second survey ran a reciprocal course and collected complementary benthic invertebrate epifauna and infauna using a benthic sled and beam trawl. The second survey provided the collections treated here. Follow-up cruises during 2007 with the same overall objectives continued the same sampling strategy along the Western Australian margin as far north as possible. Samples taken in 2007 are being identified in 2008 and will naturally add to the distributional records presented here.

Stations. Sampling was targeted at nested spatial scales of habitat – terrains of sediment and rocky substrata comprising features (mostly canyons and sediments terraces of the continental slope), within depth zones, across latitudes – to determine how biodiversity is distributed at particular scales. At the highest level, samples were allocated to enable comparison of the benthic bioregions already in use off the west and southwest coasts of Western Australia. Our collections come from 11 sites between Albany and Ningaloo (south of Exmouth) at notional depths of 100 and 400 m, and seven cross-depth transects (at intended depths of 100, 200, 400, 700 and 1000 m) made on special features of interest off Albany, Perth Canyon, Abrolhos and Ningaloo (Fig. 1). Separate targeting of hard and soft seabed terrain types was undertaken successfully in most areas.

Sampling gear. Samples were obtained using two gears, the "Sherman sled" and a beam trawl. The Sherman sled is a CMAR-designed robust sled with 1.2-metre-wide opening (0.6 m high) and is fitted with a 25 mm stretched-mesh net (Lewis, 1999). On some occasions a secondary 1 mm-mesh net was fitted inside. The beam trawl was CMAR-modified version of the French IRD design, 4 metres wide and fitted with a 25 mm stretched-mesh net.

Shipboard sorting. Samples from the sled or beam trawl were placed into one or more plastic fish boxes on deck and transferred to the wet sorting tray below deck. The material was spread out, turned and broken up and individual decapods captured and placed in 150 x 80 mm plastic dishes in seawater in rough taxonomic groups. Individuals in these dishes were further separated into operational taxonomic units (OTUs that represented our initial separation of taxa) before being labelled with provisional names and station and acquisition numbers.

Fixation and preservation. Most crustaceans were fixed in formalin but some specimens of abundant species or limbs of others were placed directly into 95% ethanol. At Museum Victoria formalin-fixed material was transferred to fresh water to soak overnight and then to 70% ethanol.

Identification. Several helpers (acknowledged below) separated the collection into more refined OTUs under the guidance of the second author who made many initial identifications. The ultimate identification of species was made by the first author with reference to general texts, in particular Poore (2004) and Sakai (1976) and the considerable primary literature cited where relevant below. Although every attempt was made to be confident of identifications no specimens were compared with types. For this reason and because so many of the determinations were of species hitherto unrecorded from Australia, the identifications must be treated as provisional.

Besides zoological names at the lowest level possible, each species was assigned a unique "MoV" number, continuing a

series started at Museum Victoria for species collected in Bass Strait in 1990. Each MoV number is permanently associated with its taxon and independent of its accepted generic or higher classification. MoV numbers are part of the taxonomy module of the Museum Victoria KEmu® registration database.

Higher taxonomy follows that of Davie (2002a; 2002b), Poore (2004) and (Ng et al., 2008) (see page 35).

Data storage. Each specimen-lot was registered on the Museum Victoria KEmu® registration catalogue from which the records published below were generated.

Presentation of taxonomic results

The results are presented as species within genera within families within infraorders (Dendrobranchiata first, Pleocyemata infraorders next in alphabetical order). The order of families is alphabetical except for Brachyura where Ng et al.'s (2008) hierarchy is followed. For each family, the species found are summarised and the literature resources used cited.

Each species is listed by name with its authority when appropriate. Uncertain identifications are prefixed “cf.” and new species close to another known species are prefixed “aff.” “MoV” numbers are given for all taxa and used as specific names for uncertain or new species.

Specimen records for each species are summarised as follows: *Records*: the total number of specimens, with latitudinal range (to nearest minute) and depth range (in metres). The latitudinal range of all samples is from 20°59'S to 35°1'S. (The longitudinal range is 112°14'E to 118°43'E.) The shallowest actual sample depth was at 50 m and the deepest at 1260 m; most measured depths are near the intended depths of 100, 400, 700 and 1000 m. A tick ✓ at the end of this line indicates that a specimen or fraction of specimen was fixed directly in alcohol (most material was fixed in formalin).

Distribution: a general comment on published distribution plus a comment on whether the species is a new species, new for southwestern Australia, new for WA or for all of Australia.

Reference: bibliographic citation used for identification.

Following the text for many species are coloured photographs. Those taken on board ship are by Karen Gowlett-Holmes. Photos of specimens taken at Museum Victoria after preservation and colour loss are by Anna McCallum or David Staples.

The entry under *Records* summarises the detailed collection data stored in the Museum Victoria KEmu® database. Sections of these data are publicly available at the Museum Victoria, Collections and Research website, “Search Natural Sciences collections”.

<http://collections.museumvictoria.com.au/browser.php?type=Zoology&phylum=Arthropoda>

Here, it is possible to search on named species and “Map Selected Specimens and Species with Google Maps”. The resulting map is derived from all Museum Victoria records, not just those from this survey. The maps reveal the bias inherent in museum collections. The map of the apparent distribution of the common slipper lobster *Ibacus alticrenatus* includes 36 records from southeastern and western Australia but none apparently from the Great Australian Bight (Fig. 2). The species certainly occurs there, as it does along a substantial part of the eastern Australian coast but no collecting by Museum Victoria has been done in these regions.

It is also possible to map the same species through an OZCAM (Online Zoological Collections of Australian Museums) query of Australia's fauna:

<http://www.ozcam.gov.au/cgi-bin/emu-dataportal.cgi>.

A search on OZCAM returns a map using collection data from all relevant Australia museums.

It is not possible to search for undescribed species. A search on a genus is likely to return results for more than one species.



Figure 2. The apparent distribution of *Ibacus alticrenatus* based on 36 records from Museum Victoria. The absence from the south coast is unlikely to be real.

Taxonomic results and commentary

The collection of ~6083 specimens representing 524 provisional species is the first comprehensive characterisation of the fauna of the continental margin of southwestern Australia. For comparison, Poore's (2004) identification guide to southern Australian marine decapods includes 800 species and the *Zoological Catalogue of Australia* enumeration of all named Australian marine Decapoda (Davie, 2002a, b) listed 2077 marine species. Poore's (2004) guide covered southern Australia extending on the west coast as far north as Perth (31°S). Our estimate is that 76 species previously unrecorded south of Perth were found in this survey, i.e., a 9.4% increase over Poore's enumeration from museum collections and literature.

The survey illustrates how little is known about the fauna of the continental margin of most of Australia. The eastern slope of NSW and Tasmania is best known. These collections are the first systematic samples from southern WA.

Overall, 175 species (33%) were new to science (Table 1). This figure is based on what we feel is a thorough review of the literature covering the fauna of Australia and the Indo-West Pacific. The number is probably an underestimate and is subject to further examination by taxonomic experts. Many of the so-called “new records” (88 species for Australia as a whole, 62 for WA and 69 for southern WA) may well prove to be new species, different from the similar species with which they have been identified. The highest percentage of new species was in Thalassinidea (83% of 23 species), much higher than the next most novel infraorders (50% of 127 species of Anomura and 31% of 227 species of Brachyura).

Many species were rare. Forty-two per cent (222 species) were found in just one of 127 samples and a further 17% (89 species) in only two samples. This is a common feature of exploration of this type and hints that the number of species yet to be discovered is much larger than anticipated.

New Australian records (88 species or 6%) were characterised as such because they did not appear in Davie's catalogues. Most were species already described from the Indo-West Pacific region (tropical and subtropical regions from Japan through to east Africa). In all cases, lack of time or few specimens prevented a thorough comparison between the WA material and original descriptions. Identifications in this category should be treated as probable at best – several may well be additional new species.

It is notable that several deep water species recently reported from Tasmanian seamounts have (with few exceptions) not been rediscovered in southern WA (Ahyong and Poore, 2004a, b).

Invitation

The process of identification of Decapoda necessitates familiarity with diverse morphologies, and access to many keys and descriptions. Most decapod taxonomists specialise in one or few families (either hermit crabs, or some crabs or prawns).

No-one is a specialist in all 77 families recognised here. These results have depended on consulting the 188 original research papers and books cited below. Poore's guidebook to southern (south of 31°S on the west coast) Australian decapods included only 24% of the species discovered in the southwest at these latitudes and a much smaller percentage of the total fauna. The collection offers considerable scope for taxonomic, evolutionary and biogeographic study. The material is available for study at Museum Victoria or on loan to crustacean taxonomists world-wide.

Table 1. Summary of numbers of species in genera, families and infraorders, including numbers of new Australian records, new records for Western Australia, and new records for southwestern Australia. Dendrobranchiata are listed first and infraorders of Pleocyemata next in alphabetical order.

Infraorder	Family	Genus	Total species	New Australian species	New WA species	New record for S WA	New species	% new spp
Dendrobranchiata	Aristeidae	<i>Aristeus</i>	4	1	1	1		0%
	Aristeidae	<i>Pseudaristeus</i>	1					0%
	Aristeidae		5	1	1	1	0	0%
	Benthescymidae	<i>Benthescymus</i>	1					0%
	Benthescymidae		1	0	0	0	0	0%
	Penaeidae	<i>Metapenaeopsis</i>	7	2	1	1		0%
	Penaeidae	<i>Parapenaeus</i>	5	4	1			0%
	Penaeidae	<i>Penaeopsis</i>	2				2	100%
	Penaeidae	<i>Penaeus</i>	1					0%
	Penaeidae	<i>Trachypenaeus</i>	1					0%
	Penaeidae		16	6	2	1	2	13%
	Sergestidae	<i>Sergestes</i>	2					0%
	Sergestidae	<i>Sergia</i>	2	1				0%
	Sergestidae		4	1	0	0	0	0%
	Sicyoniidae	<i>Sicyonia</i>	4	2			1	25%
	Sicyoniidae		4	2	0	0	1	25%
	Solenoceridae	<i>Hadropenaeus</i>	1			1		0%
	Solenoceridae	<i>Haliporoides</i>	1					0%
	Solenoceridae	<i>Haliporus</i>	1			1		0%
	Solenoceridae	<i>Hymenopenaeus</i>	2		1			0%
	Solenoceridae	<i>Solenocera</i>	9		1	5		0%
	Solenoceridae		14	0	2	7	0	0%
Dendrobranchiata	all taxa		44	10	5	9	3	7%
Achelata	Palinuridae	<i>Puerulus</i>	1			1	0	0%
	Palinuridae		1	0	0	1	0	0%
	Scyllaridae	<i>Crenarctus</i>	1				0	0%
	Scyllaridae	<i>Ibacus</i>	2				0	0%
	Scyllaridae	<i>Remiarctus</i>	1			1	0	0%
	Scyllaridae		2	0	0	1	0	0%
Achelata	all taxa		4	0	0	2	0	0%
Anomura	Chirostylidae	<i>Uroptychus</i>	5				1	20%
	Chirostylidae		5	0	0	0	1	20%
	Galatheidae	<i>Agononida</i>	6	1	2	1	2	33%
	Galatheidae	<i>Allogalathea</i>	1					0%
	Galatheidae	<i>Enriquea</i>	1		1			0%
	Galatheidae	<i>Galathea</i>	7	1	2		4	57%
	Galatheidae	<i>Lauriea</i>	1					0%
	Galatheidae	<i>Munida</i>	19	2	5		11	58%
	Galatheidae	<i>Munidopsis</i>	7	4		1		0%
	Galatheidae	<i>Paramunida</i>	1	1				0%
	Galatheidae	<i>Phylladiorhynchus</i>	1					0%
	Galatheidae	<i>Raymunida</i>	1				1	100%
	Galatheidae		45	9	10	2	18	40%
	Porcellanidae	<i>Lissoporcellana</i>	1				1	100%
	Porcellanidae	<i>Pachycheles</i>	1			1		0%
	Porcellanidae	<i>Petrolisthes</i>	2					0%

Infraorder	Family	Genus	Total species	New Australian species	New WA species	New record for S WA	New species	% new spp
Anomura (cont.)	Porcellanidae	<i>Polyonyx</i>	1					0%
	Porcellanidae	<i>Porcellanella</i>	1					0%
	Porcellanidae		6	0	0	1	1	17%
	Albuneidae	<i>Albunea</i>	1					0%
	Albuneidae	<i>Stemonopa</i>	1					0%
	Albuneidae		2	0	0	0	0	0%
	Diogenidae	<i>Calcinus</i>	4				4	100%
	Diogenidae	<i>Ciliopagurus</i>	1	1				0%
	Diogenidae	<i>Dardanus</i>	5				5	100%
	Diogenidae	<i>Diogenes</i>	1				1	100%
	Diogenidae	<i>Paguristes</i>	7		1		4	57%
	Diogenidae	<i>Strigopagurus</i>	1					0%
	Diogenidae		19	1	1	0	14	74%
	Lithodidae	<i>Lithodes</i>	1				1	100%
	Lithodidae	<i>Paralomis</i>	1				1	100%
	Lithodidae		2	0	0	0	2	100%
	Paguridae	<i>Anapagrides</i>	1				1	100%
	Paguridae	<i>Bathypaguropsis</i>	1		1			0%
	Paguridae	<i>Cestopagurus</i>	1				1	100%
	Paguridae	<i>Hemipagurus</i>	1				1	100%
	Paguridae	<i>Lophopagurus</i>	2		2			0%
	Paguridae	<i>Michelopagurus</i>	1				1	100%
	Paguridae	<i>Nematopagurus</i>	3				3	100%
	Paguridae	<i>Porcellanopagurus</i>	1	1				0%
	Paguridae	<i>Propagurus</i>	1	1				0%
	Paguridae	<i>Pylopaguropsis</i>	2				1	50%
	Paguridae	<i>Spiropagurus</i>	1		1			0%
	Paguridae	<i>Turleana</i>	2	2				0%
	Paguridae	<i>Pagurid</i>	16				16	100%
	Paguridae		33	4	4	0	24	73%
	Parapaguridae	<i>Oncopagurus</i>	3		1	2		0%
	Parapaguridae	<i>Paragiopagurus</i>	4	1			2	50%
	Parapaguridae	<i>Parapagurus</i>	1		1			0%
	Parapaguridae	<i>Strobopagurus</i>	1				1	100%
	Parapaguridae	<i>Sympagurus</i>	4		2	1		0%
	Parapaguridae		13	1	4	3	3	23%
	Pylochelidae	<i>Pylocheles</i>	1		1			0%
	Pylochelidae	<i>Pylochelidae</i>	1					0%
	Pylochelidae		2	0	1	0	0	0%
Anomura	all taxa		127	15	20	6	63	50%
Astacidea	Nephropidae	<i>Metanephrops</i>	2					0%
	Nephropidae	<i>Nephropsis</i>	2			1		0%
	Nephropidae		4	0	0	1	0	0%
	all taxa		4	0	0	1	0	0%
Brachyura	Cyclodorippidae	<i>Krangalangia</i>	1			1		0%
	Cyclodorippidae	<i>Tymolus</i>	2			2		0%
	Cyclodorippidae		3	0	0	3	0	0%
	Cymonomidae	<i>Cymonomus</i>	2	1			1	50%
	Cymonomidae		2	1	0	0	1	50%
	Dromiidae	<i>Austrodromidia</i>	1					0%

Infraorder	Family	Genus	Total species	New Australian species	New WA species	New record for S WA	New species	% new spp
Brachyura (cont.)	Dromiidae	<i>Dromia</i>	1					0%
	Dromiidae	<i>Epigodromia</i>	1				1	100%
	Dromiidae	<i>Fultodromia</i>	2				1	50%
	Dromiidae	<i>Takedromia</i>	1				1	100%
	Dromiidae		6	0	0	0	3	50%
	Dynomenidae	<i>Hirsutodynomena</i>	1			1		0%
	Dynomenidae		1	0	0	1	0	0%
	Homolidae	<i>Dagnaudus</i>	1					0%
	Homolidae	<i>Homola</i>	1					0%
	Homolidae	<i>Homologenus</i>	2	1				0%
	Homolidae	<i>Latreillopsis</i>	1					0%
	Homolidae	<i>Paramolopsis</i>	1					0%
	Homolidae	<i>Yaldwynopsis</i>	1				1	100%
	Homolidae		7	1	0	0	1	14%
	Latreilliidae	<i>Eplumula</i>	1					0%
	Latreilliidae	<i>Latreillia</i>	1	1				0%
	Latreilliidae		2	1	0	0	0	0%
	Raninidae	<i>Cosmonotus</i>	1			1		0%
	Raninidae	<i>Lyreidus</i>	2	1				0%
	Raninidae	<i>Notosceles</i>	1					0%
	Raninidae	<i>Umalia</i>	1			1		0%
	Raninidae		5	1	0	2	0	0%
	Aethridae	<i>Actaeomorpha</i>	1			1		0%
	Aethridae	<i>Drachiella</i>	1			1		0%
	Aethridae		2	0	0	2	0	0%
	Calappidae	<i>Calappa</i>	3	1	1			0%
	Calappidae	<i>Mursia</i>	3		2		1	33%
	Calappidae		6	1	3	0	1	17%
	Atelecyclidae	<i>Trichopeltarion</i>	2				2	100%
	Atelecyclidae		2	0	0	0	2	100%
	Carpiliidae	<i>Carpilius</i>	1			1		0%
	Carpiliidae		1	0	0	1	0	0%
	Corystidae	<i>Gomeza</i>	1					0%
	Corystidae	<i>Jonas</i>	1				1	100%
	Corystidae		2	0	0	0	1	50%
	Dorippidae	<i>Dorippe</i>	1					0%
	Dorippidae	<i>Neodorippe</i>	1	1				0%
	Dorippidae	<i>Paradorippe</i>	1			1		0%
	Dorippidae		3	1	0	1	0	0%
	Ethusidae	<i>Ethusa</i>	3				3	100%
	Ethusidae	<i>Ethusina</i>	1				1	100%
	Ethusidae		4	0	0	0	4	100%
	Hypothalassiidae	<i>Hypothalassia</i>	1					0%
	Hypothalassiidae		1	0	0	0	0	0%
	Chasmocarcinidae	<i>Camatopsis</i>	2	1			1	50%
	Chasmocarcinidae	<i>Megaesthesius</i>	1	1				0%
	Chasmocarcinidae		3	1	0	0	0	0%
	Euryplacidae	<i>Heteroplax</i>	2				2	100%
	Euryplacidae		2	0	0	0	2	100%
	Goneplacidae	<i>Carcinoplax</i>	2	1			2	100%
	Goneplacidae	<i>Notonyx</i>	1	1				0%
	Goneplacidae	<i>Psopheticus</i>	1	1				0%
	Goneplacidae	<i>Pycnoplax</i>	5			1	3	60%
	Goneplacidae		9	2	0	1	5	56%

Infraorder	Family	Genus	Total species	New Australian species	New WA species	New record for S WA	New species	% new spp
Brachyura (cont.)	Mathildellidae	<i>Mathildella</i>	1	1				0%
	Mathildellidae	<i>Platypilumnus</i>	1			1		0%
	Mathildellidae	<i>Mathildellid</i>	1				1	100%
	Mathildellidae		3	1	0	1	1	33%
	Hexapodidae	<i>Hexaplax</i>	1	1				0%
	Hexapodidae		1	1	0	0	0	0%
	Iphiculidae	<i>Iphiculus</i>	1			1		0%
	Iphiculidae		1	0	0	1	0	0%
	Leucosiidae	<i>Arcania</i>	8	2	2	1	2	25%
	Leucosiidae	<i>Ebalia</i>	4				3	75%
	Leucosiidae	<i>Leucosia</i>	4			1	1	25%
	Leucosiidae	<i>Merocryptus</i>	1					0%
	Leucosiidae	<i>Myra</i>	3				2	67%
	Leucosiidae	<i>Myrine</i>	1			1		0%
	Leucosiidae	<i>Oreophorus</i>	1	1				0%
	Leucosiidae	<i>Parilia</i>	1	1				0%
	Leucosiidae	<i>Philyra</i>	1				1	100%
	Leucosiidae	<i>Randallia</i>	6	1			4	67%
	Leucosiidae		30	5	2	3	13	43%
	Epialtidae	<i>Austrolibinia</i>	1			1		0%
	Epialtidae	<i>Griffinia</i>	1					0%
	Epialtidae	<i>Hyastenus</i>	1			1		0%
	Epialtidae	<i>Lahaina</i>	1					0%
	Epialtidae	<i>Naxioides</i>	3		3			0%
	Epialtidae	<i>Phalangipus</i>	2			1		0%
	Epialtidae	<i>Rochinia</i>	5	1	1		3	60%
	Epialtidae		14	1	4	3	3	21%
	Hymenosomatidae	<i>Halicarcinus</i>	1				1	100%
	Hymenosomatidae	<i>Trigonoplax</i>	1					0%
	Hymenosomatidae		2	0	0	0	1	50%
	Inachidae	<i>Achaeus</i>	5			1	1	20%
	Inachidae	<i>Camposcia</i>	1			1		0%
	Inachidae	<i>Cyrtomaia</i>	2					0%
	Inachidae	<i>Dorhynchus</i>	1					0%
	Inachidae	<i>Dumea</i>	1					0%
	Inachidae	<i>Ephippias</i>	1			1		0%
	Inachidae	<i>Grypachaeus</i>	1	1				0%
	Inachidae	<i>Oncinopus</i>	3		1		1	33%
	Inachidae	<i>Physachaeus</i>	1	1				0%
	Inachidae	<i>Platymaia</i>	2			1		0%
	Inachidae	<i>Pleistacantha</i>	1		1			0%
	Inachidae	<i>Sunipea</i>	1	1				0%
	Inachidae		20	3	2	4	2	10%
	Majidae	<i>Entomonyx</i>	2	1		1		0%
	Majidae	<i>Leptomithrax</i>	4			1	2	50%
	Majidae	<i>Maja</i>	3	3				0%
	Majidae	<i>Planotergum</i>	1					0%
	Majidae	<i>Prismatopus</i>	3	1			1	33%
	Majidae	<i>Majid</i>	1				1	100%
	Majidae		13	5	0	2	3	23%
	Palicidae	<i>Micropalicus</i>	1			1		0%
	Palicidae	<i>Neopalicus</i>	1			1		0%
	Palicidae	<i>Paliculus</i>	1		1			0%
	Palicidae	<i>Parapalicus</i>	1				1	100%

Infraorder	Family	Genus	Total species	New Australian species	New WA species	New record for S WA	New species	% new spp
Brachyura (cont.)	Palicidae	<i>Pseudopalicus</i>	1					0%
	Palicidae		5	0	1	2	1	20%
	Parthenopidae	<i>Aulacolambrus</i>	1				1	100%
	Parthenopidae	<i>Garthambrus</i>	2				2	100%
	Parthenopidae	<i>Parthenope</i>	1			1		0%
	Parthenopidae	<i>Platylambrus</i>	1		1			0%
	Parthenopidae	<i>Pseudolambrus</i>	1				1	100%
	Parthenopidae	<i>Rhinolambrus</i>	1				1	100%
	Parthenopidae	<i>Thyrolambrus</i>	1	1				0%
	Parthenopidae	<i>Parthenopid</i>	1				1	100%
	Parthenopidae		9	1	1	1	6	67%
	Pilumnidae	<i>Bathypilumnus</i>	1		1			0%
	Pilumnidae	<i>Caecopilumnus</i>	1	1				0%
	Pilumnidae	<i>Cryptolutea</i>	1			1		0%
	Pilumnidae	<i>Eumedonus</i>	1					0%
	Pilumnidae	<i>Heteropilumnus</i>	1				1	100%
	Pilumnidae	<i>Lophoplax</i>	1				1	100%
	Pilumnidae	<i>Mertonia</i>	1	1				0%
	Pilumnidae	<i>Paraselwynia</i>	1				1	100%
	Pilumnidae	<i>Pilumnopeus</i>	1				1	100%
	Pilumnidae	<i>Pilumnus</i>	11	4	1	1	6	55%
	Pilumnidae	<i>Pilumnid</i>	1					0%
	Pilumnidae		21	6	2	2	10	48%
	Portunidae	<i>Charybdis</i>	2	1	1			0%
	Portunidae	<i>Echinolatus</i>	1			1		0%
	Portunidae	<i>Libystes</i>	1		1			0%
	Portunidae	<i>Liocarcinus</i>	1					0%
	Portunidae	<i>Lissocarcinus</i>	1					0%
	Portunidae	<i>Lupocyclus</i>	3	1	1		1	33%
	Portunidae	<i>Nectocarcinus</i>	1					0%
	Portunidae	<i>Ovalipes</i>	2		2			0%
	Portunidae	<i>Parathranites</i>	2	1			1	50%
	Portunidae	<i>Portunus</i>	7	4			1	14%
	Portunidae	<i>Thalamita</i>	3		1	1		0%
	Portunidae	<i>Portunid</i>	1				1	100%
	Portunidae		25	7	6	2	4	16%
	Retroplumidae	<i>Retropluma</i>	1	1				0%
	Retroplumidae		1	1	0	0	0	0%
	Trapeziidae	<i>Quadrella</i>	1	1				0%
	Trapeziidae		1	1	0	0	0	0%
	Panopeidae	<i>Homoioplax</i>	1			1		0%
	Panopeidae		1	0	0	1	0	0%
	Xanthidae	<i>Actaea</i>	2					0%
	Xanthidae	<i>Atergatopsis</i>	1	1				0%
	Xanthidae	<i>Calvactaea</i>	1					0%
	Xanthidae	<i>Chlorodiella</i>	1					0%
	Xanthidae	<i>Demania</i>	1		1			0%
	Xanthidae	<i>Medaeus</i>	1				1	100%
	Xanthidae	<i>Monodaeus</i>	1	1				0%
	Xanthidae	<i>Nanocassiope</i>	2				2	100%
	Xanthidae	<i>Novactaea</i>	1					0%
	Xanthidae	<i>Palapedia</i>	2	1				0%
	Xanthidae	<i>Paractaea</i>	2				1	50%
	Xanthidae	<i>Paraxanthias</i>	1				1	100%

Infraorder	Family	Genus	Total species	New Australian species	New WA species	New record for S WA	New species	% new spp
	Xanthidae	<i>Paraxanthodes</i>	1	1				0%
	Xanthidae	<i>Platypodia</i>	1	1				0%
	Xanthidae		18	5	1	0	5	28%
Brachyura	all taxa		227	47	22	33	70	31%
Caridea	Alpheidae	<i>Alpheopsis</i>	3				3	100%
	Alpheidae	<i>Alpheus</i>	8				4	50%
	Alpheidae	<i>Synalpheus</i>	7					0%
	Alpheidae		18	0	0	0	7	39%
	Anchistioididae	<i>Anchistioides</i>	1			1		0%
	Anchistioididae		1	0	0	1	0	0%
	Bathypalaemonellidae	<i>Bathypalaemonella</i>	1			1		0%
	Bathypalaemonellidae		1	0	0	1	0	0%
	Bresiliidae	<i>Discias</i>	1		1			0%
	Bresiliidae		1	0	1	0	0	0%
	Campylonotidae	<i>Campylonotus</i>	1					0%
	Campylonotidae		1	0	0	0	0	0%
	Crangonidae	<i>Aegaeon</i>	1		1			0%
	Crangonidae	<i>Metacrangon</i>	1				1	100%
	Crangonidae	<i>Parapontocaris</i>	2	2				0%
	Crangonidae	<i>Parapontophilus</i>	1	1				0%
	Crangonidae	<i>Philocheras</i>	2				2	100%
	Crangonidae	<i>Pontocaris</i>	2	1	1			0%
	Crangonidae	<i>Sabinea</i>	1				1	100%
	Crangonidae		10	4	2	0	4	40%
	Eugonatonotidae	<i>Eugonatonotus</i>	1			1		0%
	Eugonatonotidae		1	0	0	1	0	0%
	Glyphocrangonidae	<i>Glyphocrangon</i>	4	1		1	1	25%
	Glyphocrangonidae		4	1	0	1	1	25%
	Hippolytidae	<i>Eualus</i>	1		1			0%
	Hippolytidae	<i>Lebbeus</i>	1				1	100%
	Hippolytidae	<i>Lysmata</i>	1			1		0%
	Hippolytidae	<i>Merhippolyte</i>	1		1			0%
	Hippolytidae	<i>Tozeuma</i>	1		1			0%
	Hippolytidae		5	0	3	1	1	20%
	Nematocarcinidae	<i>Nematocarcinus</i>	4			1	1	25%
	Nematocarcinidae		4	0	0	1	1	25%
	Oplophoridae	<i>Acanthephyra</i>	3	1		1		0%
	Oplophoridae	<i>Janicella</i>	1			1		0%
	Oplophoridae	<i>Oplophorus</i>	2					0%
	Oplophoridae	<i>Systellaspis</i>	1					0%
	Oplophoridae		7	1	0	2	0	0%
	Palaemonidae	<i>Periclimenes</i>	1	1				0%
	Palaemonidae	<i>Palaemonid</i>	1				1	100%
	Palaemonidae		2	1	0	0	1	50%
	Pandalidae	<i>Chlorotocella</i>	1					0%
	Pandalidae	<i>Chlorotocus</i>	1				1	100%
	Pandalidae	<i>Heterocarpoides</i>	1	1				0%
	Pandalidae	<i>Heterocarpus</i>	5		1	2	1	20%
	Pandalidae	<i>Plesionika</i>	12	2	2	6	2	17%
	Pandalidae	<i>Procletes</i>	1			1		0%
	Pandalidae		21	3	3	9	4	19%
	Pasiphaeidae	<i>Alainopasiphaea</i>	1					0%

Infraorder	Family	Genus	Total species	New Australian species	New WA species	New record for S WA	New species	% new spp
Caridea (cont.)	Pasiphaeidae	<i>Eupasiphae</i>	1				1	100%
	Pasiphaeidae	<i>Leptochela</i>	1		1			0%
	Pasiphaeidae	<i>Pasiphaea</i>	3		1			0%
	Pasiphaeidae		6	0	2	0	1	17%
	Processidae	<i>Hayashidonus</i>	1	1				0%
	Processidae	<i>Processa</i>	2	1	1			0%
	Processidae		3	2	1	0	0	0%
	Rhynchocinetidae	<i>Rhynchocinetes</i>	2		2			0%
	Rhynchocinetidae		2	0	2	0	0	0%
	Thalassocarididae	<i>Thalassocaris</i>	1	1				0%
	Thalassocarididae		1	1	0	0	0	0%
Caridea	all taxa		88	13	14	17	20	23%
Polychelata	Polychelidae	<i>Pentacheles</i>	1				0	0%
	Polychelidae	<i>Polycheles</i>	4	1	1		0	0%
	Polychelidae		5	1	1	0	0	0%
Polychelata	all taxa		5	1	1	0	0	0%
Stenopodidea	Stenopodidae	<i>Engystenopus</i>	1	1				0%
	Stenopodidae	<i>Odontozona</i>	1			1	0	0%
	Stenopodidae		2	1	0	1	0	0%
Stenopodidea	all taxa		2	1	0	1	0	0%
Thalassinidea	Axiidae	<i>Acanthaxius</i>	1				1	100%
	Axiidae	<i>Axiopsis</i>	2	1			1	50%
	Axiidae	<i>Bouvieraxius</i>	1				1	100%
	Axiidae	<i>Calocarides</i>	2				2	100%
	Axiidae	<i>Dorphanaxius</i>	1				1	100%
	Axiidae	<i>Marianaxius</i>	1				1	100%
	Axiidae	<i>Axiid</i>	2				2	100%
	Axiidae		10	1	0	0	9	90%
	Callianassidae	<i>Callianassa</i>	5				5	100%
	Callianassidae	<i>Corallianassa</i>	1				1	100%
	Callianassidae		6	0	0	0	6	100%
	Calocarididae	<i>Ambiaxius</i>	1				1	100%
	Calocarididae		1	0	0	0	1	100%
	Gourretiidae	<i>Lipkecallianassa</i>	1				1	100%
	Gourretiidae		1	0	0	0	1	100%
	Micheleidae	<i>Michelea</i>	1				1	100%
	Micheleidae	<i>Tethisea</i>	1				1	100%
	Micheleidae		2	0	0	0	2	100%
	Upogebiidae	<i>Upogebia</i>	3				0	0%
	Upogebiidae		3	0	0	0	0	0%
Thalassinidea	all taxa		23	1	0	0	19	83%
ALL DECAPODA			524	88	62	69	175	33%

Dendrobranchiata – prawns

The Australian fauna is diverse and well studied. Many records are of benthopelagic species. Pérez Farfante & Kensley (1997) provided keys to families and genera but this work is supplemented by family and genus treatments. Forty-four species were recorded. Ten were new records for Australia, 11 range extensions along the WA coast to known Australian species and three probable new species.

Aristeidae

Of five species one is new to Australia and one to WA (Dall, 2001).

Aristeus cf. *mabahissae* Ramadan, 1938

MoV sp. 5301

Records: 4 specimens, 29°03'S–35°31'S, 1000–1076 m

Distribution: Indo-West Pacific

Reference: figure of *A. mabahissae* from Dall (2001)

Aristeus cf. *pallicauda* Komai, 1993

MoV sp. 5320

Records: 5 specimens, 21°58'S–22°04'S, 170–387 m

Distribution: Japan; first record for Australia if this species

Reference: figure of *A. pallicauda* from Dall (2001)

Aristeus semidentatus Bate, 1881

MoV sp. 5467

Records: 2 specimens, 31°57'S, 928–1170 m

Distribution: Indo-West Pacific; first record for WA

Reference: Dall (2001) [photo below]



Aristeus virilis (Bate, 1881)

MoV sp. 5465

Records: 8 specimens, 33°02'S–35°16'S, 978–1021 m

Distribution: Indo-West Pacific; first record for S WA

Reference: Dall (2001)

Pseudaristeus sibogae (De Man, 1911)

MoV sp. 5468

Records: 3 specimens, 21°58'S–22°00'S, 726–1010 m

Distribution: Indian Ocean, S Australia

Reference: Dall (2001)

Benthescymidae

A single well-known species was recorded (Dall, 2001).

Benthescymus investigatoris Alcock & Anderson, 1899

MoV sp. 5469

Records: 3 specimens, 21°56'S–29°03'S, 1000–1056 m

Distribution: Indo-West Pacific

Reference: Dall (2001)

Penaeidae

Of 16 species identified, only five could be confidently assigned to known Australia species. Six were new records for Australia (if not new species) and two probable new species. Crosnier (1985; 1991) provided key references for the most diverse genera and Grey et al. (1983) to the larger prawns.

Metapenaeopsis crassimana Racek & Dall, 1965

MoV sp. 5479

Records: 40 specimens, 24°37'S, 100 m ✓

Distribution: N, W and S Australia

Reference: Grey et al. (1983)

Metapenaeopsis aff. *difficilis* Crosnier, 1991

MoV sp. 5460

Records: 17 specimens, 21°58'S, 107 m

Distribution: Philippines, New Caledonia; new Australian record if correctly identified

Reference: figure of *M. difficilis* from Crosnier (1991)

Metapenaeopsis aff. *vaillanti* (Nobili, 1904)

MoV sp. 5462

Records: 1 specimen, 24°37'S, 100 m

Distribution: Red Sea; new Australian record if correctly identified

Reference: figure of *M. vaillanti* from Crosnier (1991)

Metapenaeopsis rosea Racek & Dall, 1965

MoV sp. 5480

Records: 28 specimens, 20°59'S–24°37'S, 100–170 m ✓

Distribution: N and E Australia; new record for WA

Reference: Grey et al. (1983)

Metapenaeopsis velutina (Dana, 1852)

MoV sp. 5476

Records: numerous specimens, 22°50'S–27°03'S, 100 m

Distribution: Indo-West Pacific including Australia

Reference: Crosnier (1991) [photo below]



Metapenaeopsis sp. MoV 5458

MoV sp. 5458

Records: 9 specimens, 22°50'S–35°11'S, 100–402 m

Distribution: possible new species close to *M. commensalis*

Reference: Crosnier (1991)

Metapenaeopsis sp. MoV 5459

MoV sp. 5459

Records: 33 specimens, 21°59'S–22°04'S, 165–206 m

Distribution: possible new species

Reference: Crosnier (1991)

Parapenaeus fissuroides Crosnier, 1986

MoV sp. 5307

Records: 4 specimens, 21°59'S–22°04'S, 165–206 m

Distribution: Indo-West Pacific; new record for Australia

Reference: Crosnier (1985)

Parapenaeus fissurus (Bate, 1881)

MoV sp. 5478

Records: 9 specimens, 21°58'S–27°55'S, 106–253 m

Distribution: W Pacific; new record for Australia

Reference: Crosnier (1985)

Parapenaeus longipes Alcock, 1905

MoV sp. 5308

Records: 1 specimen, 22°04'S, 206–201 m

Distribution: W Pacific; new record for Australia

Reference: Crosnier (1985)

Parapenaeus murrayi Ramadan, 1938

MoV sp. 5481

Records: 24 specimens, 21°00'S–28°59'S, 324–411 m

Distribution: W Pacific; new record for Australia

Reference: Crosnier (1985)

Parapenaeus sextuberculatus Kubo, 1949

MoV sp. 5482

Records: 8 specimens, 22°04'S–24°33'S, 388–399 m

Distribution: Indo-West Pacific including E Australia; new record for WA

Reference: Crosnier (1985)

Penaeopsis sp. MoV 5466

Records: 2 specimens, 21°58'S, 356–324 m

Distribution: new species

Reference: Pérez Farfante (1980)

Penaeopsis sp. MoV 5471

Records: 15 specimens, 21°00'S–21°58'S, 373–408 m

Distribution: new species

Reference: Pérez Farfante (1980)

Penaeus (Melicertus) marginatus Randall, 1840

MoV sp. 4883

Records: 1 specimen, 21°59'S, 166 m

Distribution: Indo-West Pacific; including N Australia

Reference: Grey et al. (1983)

Trachypenaeus (*Trachysalambria*) *curvirostris* (Stimpson, 1860)

MoV sp. 5309

Records: 3 specimens, 22°04'S–27°48'S, 101–123 m ✓

Distribution: Indo-West Pacific including Australia

Reference: Grey et al. (1983)

Sergestidae

Two of the four species could not be identified because the specimens were incomplete. The third is a known Australian species and the fourth a new Australian record. Vereshchaka (2000) was consulted for *Sergia*.

Sergestes sp. MoV 5453

Records: 1 specimen, 28°57'S, 678–686 m

Distribution: incomplete specimen

Reference: Pérez Farfante and Kensley (1997)

Sergestes sp. MoV 5454

Records: 2 specimens, 28°57'S–35°31'S, 678–1110 m

Distribution: incomplete specimen

Reference: Pérez Farfante and Kensley (1997)

Sergia fulgens (Hansen, 1919)

MoV sp. 5470

Records: 7 specimens, 21°58'S, 373–732 m

Distribution: Indonesia; new Australian record

Reference: Vereshchaka (2000)

Sergia prehensilis (Bate, 1881)

MoV sp. 5311

Records: 1 specimen, 35°04'S, 379 m

Distribution: cosmopolitan

Reference: Vereshchaka (2000)

Sicyonidae

Of four species, two are new Australian records and another a probable new species (Crosnier, 2003).

Sicyonia inflexa (Kubo, 1949)

MoV sp. 5312

Records: 2 specimens, 27°55'S–28°57'S, 252–686 m

Distribution: Indo-West Pacific including N WA

Reference: one of several figures in Crosnier (2003)

Sicyonia japonica Balss, 1914

MoV sp. 5313

Records: 5 specimens, 21°58'S, 107 m

Distribution: Indo-West Pacific; new Australian record

Reference: Crosnier (2003)

Sicyonia vitulans (Kubo, 1949)

MoV sp. 5314

Records: 2 specimens, 24°37'S, 100 m

Distribution: Indo-West Pacific; new Australian record

Reference: Crosnier (2003)

Sicyonia sp. MoV 5455

MoV sp. 5455

Records: 1 specimen, 35°20'S, 213 m

Distribution: new species

Reference: Crosnier (2003)

Solenoceridae

All 14 species were identified using Dall (1999). All had been previously recorded from northern Australia but the southern or western ranges of nine were extended.

Hadropenaeus lucasii (Bate, 1881)

MoV sp. 5315

Records: numerous specimens, 21°00'S–35°10'S, 95–528 m

Distribution: Indo-West Pacific including Australia; first record for S WA

Reference: Dall (1999)

Haliporoides sibogae (De Man, 1907)

MoV sp. 5316

Records: numerous specimens, 21°58'S–27°08'S, 356–408 m

Distribution: Indo-West Pacific including Australia

Reference: Dall (1999) [photo below]



Haliporus taprobanensis Alcock & Anderson, 1899

MoV sp. 5317

Records: 2 specimens, 21°58'S, 690–732 m

Distribution: Indo-West Pacific; first record for S WA

Reference: Dall (1999)

Hymenopenaeus halli Bruce, 1966

MoV sp. 5461

Records: 2 specimens, 21°58'S–22°00'S, 373–1085 m

Distribution: Indo-West Pacific including E Australia; first record for WA

Reference: Dall (1999)

Hymenopenaeus propinquus (De Man, 1907)

MoV sp. 5319

Records: 3 specimens, 21°58'S–22°00'S, 658–754 m

Distribution: Indo-West Pacific including Australia

Reference: Dall (1999)

Solenocera annectens (Wood-Mason, 1891)

MoV sp. 5320

Records: 1 specimen, 21°57'S, 690–702 m

Distribution: Philippines, Indonesia, WA; first record for S WA

Reference: Dall (1999)

Solenocera barunajaya Crosnier, 1994

MoV sp. 5463

Records: 14 specimens, 21°58'S–29°52'S, 373–414 m

Distribution: N WA and Arafura Sea; first record for S WA

Reference: Dall (1999)

Solenocera choprai Nataraj, 1945

MoV sp. 5324

Records: 1 specimen, 22°04'S, 102 m

Distribution: Indo-West Pacific including N Australia; first record for S WA

Reference: Dall (1999)

Solenocera comata Stebbing, 1915

MoV sp. 5324

Records: 36 specimens, 24°33'S–24°33'S, 368–404 m ✓

Distribution: Indo-West Pacific including N WA; first record for S WA

Reference: Dall (1999)

Solenocera koelbeli De Man, 1911

MoV sp. 5326

Records: 1 specimen, 21°58'S, 177–170 m

Distribution: Indo-West Pacific including N WA

Reference: Dall (1999)

Solenocera melantho De Man, 1907

MoV sp. 5464

Records: 3 specimens, 21°58'S, 177–170 m

Distribution: Indo-West Pacific including N WA

Reference: Dall (1999)

Solenocera pectinata (Bate, 1880)

MoV sp. 5327

Records: 5 specimens, 20°59'S, 100 m

Distribution: Indo-West Pacific including Australia; first record for WA

Reference: Dall (1999)

Solenocera pectinulata Kubo, 1949

MoV sp. 5328

Records: numerous specimens, 21°59'S–22°04'S, 100–396 m

Distribution: Indo-West Pacific including Australia

Reference: Dall (1999)

Solenocera rathbuni Ramadan, 1938

MoV sp. 5330

Records: 34 specimens, 21°59'S–24°01'S, 100–166 m ✓

Distribution: Indo-West Pacific including N WA; first record for S WA

Reference: Dall (1999) [photo below]



Achelata – lobsters and bugs

These two families have been previously included in the Infraorder Palinura. We use the classification proposed by Ahyong and O'Meally (2004) and followed by Poore (2004).

Palinuridae

The sampling was not designed to catch lobsters but one species was taken. Its identification was confirmed with reference to Holthuis (1991).

Puerulus angulatus (Bate, 1888)

MoV sp. 4972

Records: 6 specimens, 21°58'S–22°50'S, 324–430 m ✓

Distribution: Indo-West Pacific including N Australia; new record for S WA

References: Holthuis (1991); Griffin & Stoddart (1995) [photo below]



Scyllaridae

Two species of commercially-important bugs (*Ibacus* spp.) and two of smaller scyllarids were recorded, all identifiable from Holthuis (1985; 2002) and Poore (2004).

Crenarctus crenatus (Whitelegge, 1900)

MoV sp. 4974

Records: 1 specimen, 35°10'S, 99 m

Distribution: S Australia

Reference: Holthuis (2002) [photo below]



Ibacus alticrenatus Bate, 1888

MoV sp. 3873

Records: many specimens, 21°58'S–35°04'S, 324–490 m ✓

Distribution: S Australia, common

References: Holthuis (1985; 2002) [photos below]



Ibacus peronii Leach, 1815

MoV sp. 1771

Records: 1 specimen, 24°01.43'S, 100 m

Distribution: S Australia

Reference: Poore (2004)

Remiarctus bertholdii (Paulson, 1875)

MoV sp. 4976

Records: 16 specimens, 20°59'S–22°04'S, 100–166 m ✓

Distribution: Indo-West Pacific including N Australia; new record for S WA

Reference: Holthuis (2002) [photo below]



Anomura – Hermit crabs, stone crabs, frog crabs and squat lobsters

Families of this diverse group are listed in three superfamilies, Galatheoidea, Hippoidea and Paguroidea. Species number 127.

Superfamily Galatheoidea

Three families were represented by 56 species. Twenty (36%) are certain or probably new species. Nine are new records for Australia of species previously reported for the Indo-West Pacific and 13 new for WA or more southern records of WA species.

Chirostylidae

Five species were separated using Ah Yong and Poore (2004a) and Baba (2005). One is a probable new species.

Uroptychus australis (Henderson, 1885)

MoV sp. 5249

Records: 4 specimens, 22°00'S–35°26'S, 658–988 m

Distribution: New Zealand, Indonesia, E Australia; first record for WA

Reference: Ah Yong and Poore (2004a) [photo below]



Uroptychus flindersi Ah Yong & Poore, 2004

MoV sp. 5447

Records: 2 specimens, 35°12'S, 431–408 m

Distribution: S Australia

Reference: Ah Yong and Poore (2004a) [photo below]



Uroptychus gracilimanus (Henderson, 1885)

MoV sp. 5248

Records: 10 specimens, 33°00'S, 397–421 m

Distribution: Indo-West Pacific including E Australia; first record for WA

Reference: Ah Yong and Poore (2004a) [photo below]



Uroptychus hesperius Ah Yong & Poore, 2004

MoV sp. 5206

Records: 1 specimen, 35°26'S, 915 m

Distribution: S WA

Reference: Ah Yong and Poore (2004a)

Uroptychus sp. MoV 5181

Records: 5 specimens, 27°48'S–29°52'S, 401–431 m

Distribution: new species

Reference: Ah Yong and Poore (2004a) [photo below]



Galatheidae

Forty-five species were represented, of which nine are new records for Australia, ten new for Western Australia and two reported more further south than previously known. Eighteen (40%) are probable new species. Baba (1988; 2005) and Ahyong and Poore (2004b) were the most relevant sources. The number of new species could well be higher if the new range extensions of Indo-West Pacific species are discovered to be new species. The genus *Munida* was richest in species (19 species) *Galathea* and *Munidopsis* with seven species each and *Agononida* with six species.

Agononida eminens (Baba, 1988)

MoV sp. 5201

Records: 8 specimens, 21°58'S–22°00'S, 658–754 m ✓

Distribution: West Pacific including E Australia; first record for WA

Reference: Baba (2005) [photo below]



Agononida incerta (Henderson, 1888)

MoV sp. 5260

Records: many specimens, 21°58'S–31°55'S, 324–754 m ✓

Distribution: Indo-West Pacific including N Australia; first record for S WA

Reference: Ahyong and Poore (2004b) [photo below]



Agononida pilosimanus (Baba, 1969)

MoV sp. 5208

Records: 9 specimens, 27°08'S–31°59'S, 414–508 m

Distribution: West Pacific including Qld; first record for WA

Reference: Baba (2005)

Agononida similis (Baba, 1988)

MoV sp. 5205

Records: 2 specimens, 21°58'S, 382 m

Distribution: Philippines; first record for Australia

Reference: Baba (2005)

Agononida sp. aff. *incerta* (Henderson, 1888)

MoV sp. 5207

Records: 5 specimens, 21°00'S–22°50.48'S, 399–430 m ✓

Distribution: new species

Reference: Ahyong and Poore (2004b)

Agononida sp. aff. *sabatesae* (Macpherson, 1994)

MoV sp. 5218

Records: 8 specimens, 31°37'S–31°59'S, 364–508 m

Distribution: new species close to New Caledonian species

Reference: Macpherson (1994)


Allogalathea elegans (Adams & White, 1848)

MoV sp. 5350

Records: 2 specimens, 22°04'S–24°37'S, 100–102 m

Distribution: Indo-West Pacific including Australia

Reference: Poore (2004: 231) [photo below]



Enriquea leviantennata (Baba, 1988)

MoV sp. 5202

Records: 3 specimens, 21°58'S–22°04'S, 373–391 m

Distribution: Indonesia, N and NE Australia; first record for WA

Reference: Baba (1988; 2005) [photo below]



Galathea aff. *orientalis* Stimpson, 1858

MoV sp. 5182

Records: 10 specimens, 22°50'S–27°03'S, 100–106 m

Distribution: new species; keys to *G. orientalis* but record for WA doubted by Baba

Reference: Haig (1974); Baba (2005)

Galathea amboinensis De Man, 1888

MoV sp. 5258

Records: 4 specimens, 24°01'S–27°48'S, 96–101 m

Distribution: Indonesia, N Qld; first record for WA

Reference: Baba (1988) [photo below]



Galathea balssi Miyake & Baba, 1964

MoV sp. 5273

Records: 3 specimens, 20°59'S–22°37'S, 100–382 m

Distribution: West Pacific including Qld; first record for WA

Reference: Baba (2005)

Galathea consobrina De Man, 1902

MoV sp. 5257

Records: 1 specimen, 34°53'S, 100–95 m ✓

Distribution: Philippines; first record for Australia

Reference: Baba (1988)

***Galathea* sp. MoV 5179**

Records: 19 specimens, 20°59'S, 100 m

Distribution: similar to *G. multilineata* from Japan–Philippines

Reference: Baba (2005)

***Galathea* sp. MoV 5209**

Records: 2 specimens, 24°01'S, 100 m

Distribution: new species

Reference: Baba (2005)

***Galathea* sp. MoV 5244**

Records: 1 specimen, 22°50'S, 100 m

Distribution: new species

Reference: Baba (2005)

Lauriea gardineri (Laurie, 1926)

MoV sp. 5259

Records: 2 specimens, 22°50'S–27°03'S, 100–382 m

Distribution: Indo-West Pacific including WA

Reference: Baba (2005), Osawa and Okuno (2004)

Munida andamanica Alcock, 1894

MoV sp. 5212

Records: 1 specimen, 21°00'S, 399–408 m

Distribution: Indo-West Pacific including N Qld; first record for WA

Reference: Baba (2005)

Munida aprosoma Ah Yong & Poore, 2004

MoV sp. 5197

Records: 8 specimens, 29°00'S–31°57'S, 700–1170 m

Distribution: NE Australia; first record for WA

Reference: Ah Yong and Poore (2004b) [photo below]



Munida babai Tirmizi & Javed, 1976

MoV sp. 5178

Records: 12 specimens, 20°59'S–21°59'S, 100–177 m ✓

Distribution: South Africa–Malaysia; first record for Australia

Reference: Baba (1988), Tirmizi and Javed (1976)

Munida disgrega Baba, 2005

MoV sp. 5210

Records: 37 specimens, 35°22'S–35°22'S, 408–680 m

Distribution: SE Australia; first record for WA

Reference: Baba (2005)

Munida haswelli Henderson, 1885

MoV sp. 3859

Records: many specimens, 21°59'S–35°11'S, 130–728 m ✓

Distribution: S Australia

Reference: Poore (2004) [photo below]



Munida heteracantha Ortmann, 1892

MoV sp. 5196

Records: 3 specimens, 21°58'S, 177–170 m

Distribution: Indo-West Pacific including Qld; first record for WA

Reference: Baba (1988) as *M. exigua*

Munida roshanei Tirmizi, 1966

MoV sp. 5180

Records: 31 specimens, 20°59'S–27°48'S, 93–123 m ✓

Distribution: Aden–Andaman Sea; first record for Australia

Reference: Baba (1988)

Munida rubridigitalis Baba, 1994

MoV sp. 5211

Records: 9 specimens, 21°00'S–24°33'S, 396–411 m ✓

Distribution: N Qld; first record for WA

Reference: Baba (1994) [photo below]



Munida* aff. *amathea Macpherson, 1995

MoV sp. 5203

Records: 1 specimen, 27°48'S, 431–416 m

Distribution: probably new species like *M. amathea* from Tuamotu

Reference: Baba (2005)

Munida* aff. *rubiesi Macpherson, 1991

MoV sp. 5183

Records: 23 specimens, 27°56'S–31°36'S, 329–704 m

Distribution: probably new species like *M. rubiesi* from Gulf of Aden

Reference: Baba (2005) [photo below]



Munida* aff. *volantis Macpherson, 2004

MoV sp. 5204

Records: 12 specimens, 27°55'S–31°55'S, 180–232 m

Distribution: probably new species like *M. volantis* from Fiji

Reference: Baba (2005) [photo below]



***Munida* sp. MoV 5176**

Records: 3 specimens, 20°59'S, 101–100 m

Distribution: probably new species like *M. janetae* from E Indian Ocean

Reference: Baba (2005)

***Munida* sp. MoV 5199**

Records: 1 specimen, 35°26'S, 912–922 m

Distribution: probably new species like *M. nesiotes* from Seychelles

Reference: Baba (2005) [photo below]



***Munida* sp. MoV 5200**

Records: 1 specimen, 33°00'S, 421–414 m

Distribution: probably new species like *M. semoni* from West Pacific

Reference: Baba (2005) [photo next page]



***Munida* sp. MoV 5214**

Records: 20 specimens, 22°04'S, 206–201 m ✓

Distribution: new species like *M. babai* from South Africa–Malaysia

Reference: Baba (2005)

***Munida* sp. MoV 5215**

Records: 2 specimens, 21°58'S, 356–324 m

Distribution: new species like *M. shaula* from W Indian Ocean

Reference: Baba (2005)

***Munida* sp. MoV 5217**

Records: 1 specimen, 22°04'S, 399–387 m

Distribution: new species

Reference: Baba (2005)

***Munida* sp. MoV 5245**

MoV sp. 5245

Records: 1 specimens, 27°03'S, 106–106 m

Distribution: new species, incomplete specimen

Reference: Baba (2005)

***Munida* sp. MoV 5526**

Records: 1 specimen, 22°004'S, 658–754 m

Distribution: new species near *M. andamanica*

Reference: Baba (2005)

***Munidopsis andamanica* MacGilchrist, 1905**

MoV sp. 5253

Records: 8 specimens, 21°58'S, 726–732 m

Distribution: West Pacific, Indonesia; first record for Australia

Reference: Baba (2005: 284) [photo below]



***Munidopsis crenatirostris* Baba, 1988**

MoV sp. 5251

Records: 17 specimens, 21°00'S–35°12'S, 396–754 m ✓

Distribution: Philippines; first record for Australia

Reference: Baba (2005) [photo below]



***Munidopsis cylindrophthalma* (Alcock, 1894)**

MoV sp. 5255

Records: 1 specimen, 21°58'S, 726–732 m

Distribution: Indo-West Pacific; first record for Australia

Reference: Baba (2005), Macpherson (2007) for colour photo

***Munidopsis dasypus* Alcock, 1894**

MoV sp. 5252

Records: 4 specimens, 29°03'S, 1000–1037 m

Distribution: Indo-West Pacific including N WA; first record for S WA

Reference: Baba and Poore (2002: 50, WA record) [photo below]



***Munidopsis kensleyi* Ahyong & Poore, 2004**

MoV sp. 5254

Records: 1 specimens, 21°55'S, 1260–1295 m

Distribution: S Australia

Reference: Baba and Poore (2002: as *M. dasypus*), Ahyong & Baba (Ahyong and Poore, 2004c)

***Munidopsis levis* (Alcock & Anderson, 1894)**

MoV sp. 5256

Records: 1 specimen, 21°58'S, 726–732 m ✓

Distribution: Andaman Sea, Philippines; first record for Australia

Reference: Baba (2005)

Munidopsis serricornis (Lovén, 1852)

MoV sp. 2677

Records: 1 specimen, 35°26'S, 900–915 m

Distribution: Indo-West Pacific including S Australia

Reference: Baba (2005), Baba and Poore (2002)

Paramunida stichas Macpherson, 1993

MoV sp. 5213

Records: 11 specimens, 23°59'S–24°33'S, 388–404 m

Distribution: Indonesia, New Caledonia; first record for Australia

Reference: Macpherson (1993)

Phylladiorhynchus pusillus (Henderson, 1885)

MoV sp. 0091

Records: 31 specimens, 23°59'S–35°10'S, 95–439 m

Distribution: Indo-West Pacific including S Australia

Reference: Poore (2004) [photo below]



Raymunida sp. MoV 5189

MoV sp. 5189

Records: 2 specimens, 21°57'S–29°48'S, 104–114 m

Distribution: new species

Reference: Macpherson and Machordom (2001) [photo below]



Porcellanidae

Five species were identified using Haig (1965) and an update (Haig, 1981). Only one was problematic, a species previously recorded from WA but possibly misidentified. Another was reported further south than previously known.

Lissoporcellana* aff. *quadrilobata (Miers, 1884)

MoV sp. 5226

Records: 12 specimens, 20°59'S, 101–100 m ✓

Distribution: probable new species like *L. quadrilobata*

Reference: Haig (1981)

Pachycheles sculptus (Milne Edwards, 1837)

MoV sp. 5221

Records: 3 specimens, 24°01'S–24°37'S, 101–100 m

Distribution: N WA; first record for S WA

Reference: Haig (1965) [photo below]



Petrolisthes militaris (Heller, 1862)

MoV sp. 5224

Records: 53 specimens, 21°59'S–28°59'S, 100–183 m ✓

Distribution: Indo-West Pacific including SW Australia

Reference: Haig (1965) [photo below]



Petrolisthes scabriculus (Dana, 1852)

MoV sp. 5220

Records: 2 specimens, 27°48.48'S, 96–98 m

Distribution: Indo-West Pacific including SW Australia

Reference: Haig (1965)

Polyonyx biunguiculatus (Dana, 1852)

MoV sp. 5225

Records: 26 specimens, 21°59'S–33°2'S, 95–166 m

Distribution: Indo-West Pacific including WA

Reference: Haig (1965) [photo below]



Porcellanella triloba White, 1851

MoV sp. 5246

Records: 1 specimen, 22°02'S, 106 m

Distribution: WA

Reference: Haig (1974)

Superfamily Hippoidea

A single family can be reported, with two species.

Albuneidae

Two species previously reported from WA were collected and identified using Boyko (2002).

Albunea oculatus Boyko, 2002

MoV sp. 5223

Records: 1 specimen, 25°54'S, 100 m

Distribution: WA

Reference: Boyko (2002: 315) [photo below]



Stemonopa insignis Efford & Haig, 1968

MoV sp. 5222

Records: 4 specimens, 24°01'S–25°54'S, 100 m ✓

Distribution: WA

Reference: Boyko (2002: 224) [photo below]



Superfamily Paguroidea

Hermit crabs are notoriously difficult to identify. Although many species have been described the literature is extremely scattered. McLaughlin (2003) has provided keys to families and genera that enabled access to the recent literature. Five families were represented among the 70 species recognised. It is highly probable that a more experienced taxonomist could identify some of the species to a lower level but the absence of both sexes proved an impediment in the many cases of rare taxa.

Diogenidae


Of 19 species, five were identifiable to species and the rest to genus level. One was a new record for Australia and another for WA. None of the recent literature reviewed by Davie (2002) or Poore (2004) proved useful. If this taxonomy is fair, three-quarters of the species discovered are new species.

Calcinus sp. MoV 5268

Records: 1 specimen, 24°37'S, 100 m

Distribution: probable new species

Reference: Morgan (1991: key) [photos below]



Calcinus sp. MoV 5389

Records: 8 specimens, 24°37'S–34°53'S, 95–100 m

Distribution: keys to *C. tropidomanus* Lewinsohn, 1981; new species or new Australian record

Reference: Poupin and McLaughlin (1998) [photo below]



Calcinus sp. MoV 5393

Records: 1 specimen, 27°48'S, 98 m

Distribution: probable new species

Reference: Poupin and McLaughlin (1998)

Calcinus sp. MoV 5396

Records: 1 specimen, 22°04'S, 102 m

Distribution: new species

Reference: Poupin and McLaughlin (1998)

Ciliopagurus cf. *krempfi* (Forest, 1952)

MoV sp. 5275

Records: 2 specimens, 22°50'S–29°48'S, 85–100 m

Distribution: new species or new Australian record; difficult to identify from key

Reference: Forest (1995: key) [photo below]



Dardanus sp. MoV 5262

Records: 2 specimens, 22°37'S–35°21'S, 92–382 m

Distribution: probable new species [photo below]



Dardanus sp. MoV 5264

Records: 1 specimen, 22°04'S, 106–101 m

Distribution: probable new species

***Dardanus* sp. MoV 5265**

Records: 2 specimens, 21°59'S, 166 m

Distribution: probable new species [photos below]



***Dardanus* sp. MoV 5266**

Records: 6 specimens, 25°55'S–33°58'S, 96–123 m

Distribution: probable new species [photo below]



***Dardanus* sp. MoV 5267**

Records: 7 specimens, 21°59'S–24°37'S, 100–166 m

Distribution: probable new species [photos below]



***Diogenes* sp. MoV 5401**

Records: 1 specimen, 24°37'S, 100 m

Distribution: probable new species

Reference: Morgan and Forest (1991)

***Paguristes aciculus* Grant, 1905**

MoV sp. 5279

Records: numerous specimens, 21°58'S–35°22'S, 100–508 m

Distribution: E Australia; new record for WA

Reference: Poore (2004: key) [photo below]



***Paguristes longisetosus* Morgan, 1987**

MoV sp. 5382

Records: 1 specimen, 22°37.04'S, 355–382 m

Distribution: S WA

Reference: Poore (2004: key) [photos below]



***Paguristes purpureantennatus* Morgan, 1987**

MoV sp. 5331

Records: 2 specimens, 31°37'S–35°10'S, 97–210 m

Distribution: S WA

Reference: Poore (2004: key) [photo below]




***Paguristes* sp. MoV 5263**

Records: 8 specimens, 21°58'S–22°02'S, 104–144 m

Distribution: new species

References: Poore (2004: key), Rahayu (2006) [photos below]



***Paguristes* sp. MoV 5277**

Records: 30 specimens, 21°00'S–22°04'S, 165–411 m

Distribution: new species

References: Poore (2004: key), Rahayu (2006) [photos below]



***Paguristes* sp. MoV 5278**

Records: 1 specimen, 21°59'S, 166 m

Distribution: new species

References: Poore (2004: key), Rahayu (2006) [photos below]



***Paguristes* sp. MoV 5394**

Records: 5 specimens, 24°33'S–27°48'S, 112–388 m

Distribution: new species

References: Poore (2004: key), Rahayu (2006) [photo below]



***Strigopagurus elongatus* Forest, 1995**

MoV sp. 1707

Records: 24 specimens, 31°37'S–35°21'S, 95–210 m

Distribution: S WA

References: Poore (2004: key) [photo below]



Lithodidae

One individual each of two species previously recorded from Tasmanian seamounts were found (Poore, 2004). Both are new species similar to named species from Japan and Peru (S. Ahyong, pers. comm.).

Lithodes aff. *longispina* Sakai, 1971

MoV sp. 2718

Records: 2 specimens, 31°58'S–35°26'S, 848–1050 m

Distribution: SE Australia; new record for WA (not *L. longispina* from Japan)

Reference: Poore (2004: 268) [photo below]



Paralomis cf. *phrixa* Macpherson, 1992

MoV sp. 2717

Records: 1 specimen, 35°26'S, 900–915 m

Distribution: Tas. Seamounts; new record for WA (not *P. phrixa* from Peru)

Reference: Poore (2004: 269) [photo below]



Paguridae

Half of all hermit crabs belong in this family. Half of the 33 species taken could not be identified beyond family level because each was represented by few specimens of only one sex. McLaughlin's (1997) work on Indonesian species includes some of those identified to species level, including a new record for Australia and another for WA. At least three-quarters (24 species) are probable new species. No one genus was especially diverse.

Anapagrises sp. MoV 5399

Records: 2 specimens, 22°04'S–31°43'S, 102 m

Distribution: females only; new record for genus in Australia

Reference: McLaughlin (2003: key to genera)

Bathypaguropsis yaldwyni McLaughlin, 1994

MoV sp. 2686

Records: 1 specimen, 31°55'S, 479–484 m

Distribution: New Zealand, Vic., Tas. Seamounts; new record for WA

Reference: McLaughlin (1994) [photo below]



Cestopagurus sp. MoV 5269

Records: 1 specimen, 31°55'S, 479–484 m

Distribution: female only; new record for genus in Australia

Reference: McLaughlin (2003: key to genera) [photo below]




Hemipagurus sp. MoV 5281

Records: 1 specimen, 22°50'S, 100 m

Distribution: probable new species; new record for genus in Australia

Reference: Asakura (2001) [photos below]



Lophopagurus (Lophopagurus) nanus (Henderson, 1888)

MoV sp. 1591

Records: 4 specimens, 31°43'S–35°20'S, 97–213 m

Distribution: S Australia; first record for WA

Reference: Poore (2004: 274) [photo below]



Lophopagurus (Australeremus) triserratus (Ortmann, 1892)

MoV sp. 5332

Records: 14 specimens, 27°03'S–35°20'S, 97–213 m

Distribution: S Australia; first record for WA

Reference: Poore (2004: 274) [photo below]



Michelopagurus sp. MoV 5280

Records: 1 specimen, 31°57'S, 928–1170 m

Distribution: first record of genus from Australia

Reference: McLaughlin (1997: 481) [photo upper right]



Nematopagurus sp. MoV 5380

Records: 2 specimens, 22°50'S, 100 m

Distribution: male only; probable new species

Reference: McLaughlin (2004) [photo below]



Nematopagurus sp. MoV 5383

Records: 2 specimens, 31°55'S–35°22'S, 194–232 m

Distribution: females only; probable new species

Reference: McLaughlin (2004) [photo below]



Nematopagurus sp. MoV 5384

Records: 15 specimens, 21°59'S, 166 m

Distribution: males and females; probable new species

Reference: McLaughlin (2004)

Porcellanopagurus filholi de Saint Laurent & McLaughlin, 2000

MoV sp. 5398

Records: 1 specimen, 29°52'S, 414–401 m

Distribution: South Africa, New Zealand; first record for Australia

Reference: McLaughlin (2000) [photo below]



Propagurus haigae (McLaughlin, 1997)

MoV sp. 5333

Records: 9 specimens, 27°48'S–35°22'S, 394–428 m

Distribution: E Australia, Indonesia; first record for WA

Reference: McLaughlin and de Saint Laurent (1998) [photo below]



Pylopaguropsis zebra (Henderson, 1893)

MoV sp. 5334

Records: 9 specimens, 21°59'S–24°37'S, 100–166 m

Distribution: Indo-West Pacific including N WA

Reference: McLaughlin and Haig (1989) [photo upper right]



Pylopaguropsis sp. MoV 5276

Records: 1 specimen, 25°54'S, 100–95 m

Distribution: male; new species

Reference: McLaughlin and Haig (1989)

Spiropagurus fimbriatus Lewinsohn, 1982

MoV sp. 5335

Records: 5 specimens, 21°58'S–22°04'S, 101–166 m

Distribution: Red Sea, N Qld; first record for WA

Reference: Lewinsohn (1982) [photo below]



Turleana albatrossae (McLaughlin & Haig, 1996)

MoV sp. 5284

Records: 8 specimens, 20°59'S–27°48'S, 96–106 m

Distribution: new record for Australia

Reference: McLaughlin and Haig (1996) [photo next page]



Turleana multispina McLaughlin, 1997

MoV sp. 5400

Records: 2 specimens, 23°59'S, 411 m

Distribution: Indonesia; new record for Australia

Reference: McLaughlin (1997)

Pagurid sp. MoV 5261

Records: 6 specimens, 22°51'S–22°50'S, 100–106 m

Distribution: can not be keyed to genus

Reference: McLaughlin (2003) [photo below]



Pagurid sp. MoV 5270

Records: 1 specimen, 29°03'S, 1000–1037 m

Distribution: male, similar to *Tomopaguropsis*

Reference: McLaughlin (2003) [photo upper right]



Pagurid sp. MoV 5271

Records: 3 specimens, 29°03'S, 1000–1037 m

Distribution: females only; can not be keyed to genus

Reference: McLaughlin (2003) [photo below]



Pagurid sp. MoV 5274

Records: 2 specimens, 35°25'S, 925–913 m

Distribution: females only; like *Lophopagurus*

Reference: McLaughlin (2003) [photo below]



***Pagurid* sp. MoV 5283**

Records: 2 specimens, 21°56'S–29°03'S, 1000–1037 m

Distribution: males only; can not be keyed to genus

Reference: McLaughlin (2003)

***Pagurid* sp. MoV 5285**

Records: 1 specimen, 35°26'S, 900–915 m

Distribution: male only; can not be keyed to genus

Reference: McLaughlin (2003)

***Pagurid* sp. MoV 5286**

Records: 2 specimens, 20°59'S, 100 m

Distribution: males only; can not be keyed to genus

Reference: McLaughlin (2003)

***Pagurid* sp. MoV 5381**

Records: 2 specimens, 21°58'S–28°59'S, 170–183 m

Distribution: females only; can not be keyed to genus

Reference: McLaughlin (2003)

***Pagurid* sp. MoV 5385**

Records: 3 specimens, 23°59'S, 398–402 m

Distribution: males only; can not be keyed to genus

Reference: McLaughlin (2003)

***Pagurid* sp. MoV 5386**

Records: 1 specimen, 22°00'S, 1085–1077 m

Distribution: female only; cannot be keyed to genus

Reference: McLaughlin (2003)

***Pagurid* sp. MoV 5387**

Records: 3 specimens, 31°58'S–31°57'S, 848–1170 m

Distribution: keys to *Parapagurodes*

Reference: McLaughlin (2003)

***Pagurid* sp. MoV 5388**

Records: 4 specimens, 31°58'S, 848–1050 m

Distribution: females only; can not be keyed to genus

Reference: McLaughlin (2003)

***Pagurid* sp. MoV 5390**

Records: 1 specimen, 22°50'S, 100 m

Distribution: male only; can not be keyed to genus

Reference: McLaughlin (2003)

***Pagurid* sp. MoV 5391**

Records: 4 specimens, 21°58'S, 107 m

Distribution: males and females; can not be keyed to genus

Reference: McLaughlin (2003)

***Pagurid* sp. MoV 5392**

Records: 2 specimens, 21°58'S–24°01'S, 100–107 m

Distribution: male and female; can not be keyed to genus

Reference: McLaughlin (2003)

***Pagurid* sp. MoV 5402**

Records: 3 specimens, 21°58'S, 732 m

Distribution: females only; can not be keyed to genus

Reference: McLaughlin (2003)

Parapaguridae

Ten of the 13 species taken were identifiable using the works of Lemaitre (1996; 2004a; 2004b). Four were new records for WA.

***Oncopagurus indicus* (Alcock, 1905)**

MoV sp. 5336

Records: 33 specimens, 21°58'S–35°25'S, 373–1037 m

Distribution: Indo-West Pacific including N Australia; new record for S WA

Reference: Lemaitre (1996) [photo below]



***Oncopagurus minutus* (Henderson, 1896)**

MoV sp. 5337

Records: 6 specimens, 21°58'S–31°58'S, 732–1050 m ✓

Distribution: Indo-West Pacific including E Australia; new record for WA

Reference: Lemaitre (1996)

***Oncopagurus monstrosus* (Alcock, 1894)**

MoV sp. 5338

Records: many specimens, 22°50'S–35°26'S, 329–1050 m ✓

Distribution: Indo-West Pacific including N WA; new record for S WA

Reference: Lemaitre (1996) [photo below]



Paragiopagurus boletifer (de Saint Laurent, 1972)

MoV sp. 5339

Records: 1 specimen, 22°37'S, 355–382 m

Distribution: Indo-West Pacific; new record for Australia

Reference: Lemaitre (1996) [photo below]



Paragiopagurus diogenes (Whitelegge, 1900)

MoV sp. 5340

Records: 36 specimens, 24°01'S–33°58'S, 96–407 m

Distribution: Indo-West Pacific including Australia

Reference: Lemaitre (1996) [photo below]



***Paragiopagurus* sp. MoV 5272**

Records: 25 specimens, 21°00'S–33°00'S, 355–1010 m

Distribution: new species

Reference: Lemaitre (1996) [photo below]



***Paragiopagurus* sp. MoV 5397**

Records: 9 specimens, 24°33'S, 388–368 m

Distribution: new species

Reference: Lemaitre (1996)

Parapagurus latimanus Henderson, 1888

MoV sp. 5341

Records: 26 specimens, 22°00'S–35°23'S, 479–1110 m

Distribution: Indo-West Pacific including E Australia; first record for WA

Reference: Lemaitre (1999) [photo below]




***Strobopagurus* sp. MoV 5282**

Records: 11 specimens, 21°58'S–31°37'S, 364–1037 m ✓

Distribution: new species

Reference: Lemaitre (2004b) [photo next page]




Sympagurus brevipes (de Saint Laurent, 1972)

MoV sp. 5342

Records: 1 specimen, 21°58'S, 726–732 m

Distribution: Indo-West Pacific including N Australia; new record for S WA

Reference: Lemaitre (1996) [photos below]



Sympagurus dimorphus (Studer, 1883)

MoV sp. 5343

Records: 29 specimens, 31°55'S–35°22'S, 423–680 m

Distribution: Southern Ocean including Australia; first record for WA

Reference: Lemaitre (1996) [photo below]



Sympagurus planimanus (de Saint Laurent, 1972)

MoV sp. 5344

Records: 71 specimens, 21°58'S, 726–732 m

Distribution: West Pacific including N WA

Reference: Lemaitre (1996)

Sympagurus villosus Lemaitre, 1996

MoV sp. 5345

Records: 2 specimens, 21°58'S–22°00'S, 324–1010 m

Distribution: eastern Australia; first record for WA

References: Lemaitre (1996), Poore (2004)

Pylochelidae

No pylochelids have been recorded for WA but the only identified species ranges across the Indian and SW Pacific. The second was represented by juveniles. The systematics of the family was reviewed by Forest (1987).

Pylocheles mortensenii Boas, 1926

MoV sp. 5346

Records: 1 specimen, 31°37'S, 364–404 m

Distribution: Indo-West Pacific including Qld, New Zealand; first record for WA

Reference: Forest (1987) [photos below]



Pylochelidae sp. MoV 5395

Records: 2 specimens, 24°33'S, 388–368 m

Distribution: juvenile specimens

Reference: Forest (1987)

Astacidea – scampi

Astacidea are represented in these collections by one family that includes some species of commercial interest.

Nephropidae

Four well-known species in two genera were recorded, all identifiable from Poore (2004) or Macpherson (1990; 1993). One is a new record for southern WA. The papers cited have figures.

Metanephrops boschmai (Holthuis, 1964)

MoV sp. 5067

Records: 17 specimens, 21°58'S–35°13'S, 324–554 m ✓

Distribution: S and W Australia

Reference: Poore (2004: 165) [photos below]



Metanephrops velutinus Chan & Y3u, 1991

MoV sp. 5077

Records: 13 specimens, 22°04'S–35°12'S, 387–508 m

Distribution: West Pacific including Australia

Reference: Poore (2004: 165) [photo below]



Nephropsis acanthura Macpherson, 1990

MoV sp. 4968

Records: 2 specimens, 21°55'S, 1260–1295 m ✓

Distribution: Indo-West Pacific including S Australia

Reference: Poore (2004: 166)

Nephropsis stewarti Wood-3Mason, 1872

MoV sp. 5068

Records: 1 specimen, 31°58'S, 848–1050 m

Distribution: Indo-West Pacific including N Australia; new record for S WA

Reference: Macpherson (1993) [photos below]



Brachyura – crabs

Thirty-two families were represented by 227 nominal species. The single reference to brachyuran crabs from a broad geographic region of Australia (Poore, 2004) was found to deal with only a small fraction of the species found. Numerous papers, especially recent works describing species from the Western Pacific and Indonesia, were consulted to make species determinations.

One quarter of all species (47 species) were first records for Australia, a further 22 species first records for WA and 31 first records for southern WA. Seventy-one species (31%) are new species. Some of the species noted as new to Australia or to WA should be considered tentative identifications until specimens are compared with types or representatives from type localities.

The family arrangement and sequence is that of Ng et al. (2008). Their list of all known species updates the classification used by Poore (2004). Families are listed in this sequence, genera and species alphabetically within families. The eubrachyuran subsection Thoracotremata was not represented.

Section Podotremata

Superfamily Cyclodorippoidea

Cyclodorippidae

Cymonomidae

Superfamily Dromoidea

Dromiidae

Dynomenidae

Superfamily Homoloidea

Homolidae

Latreilliidae

Superfamily Raninoidea

Raninidae

Section Eubrachyura

Subsection Heterotremata

Superfamily Aethroidea

Aethridae

Superfamily Calappoidea

Calappidae

Superfamily Cancroidea

Atelecyclidae

Superfamily Carpiloidea

Carpiliidae

Superfamily Corystoidea

Corystidae

Superfamily Dorippoidea

Dorippidae

Ethusidae

Superfamily Eriphioidea

Hypothalassiidae

Superfamily Goneplacoidea

Chasmacardinidae

Euryplacidae

Goneplacidae

Mathildellidae

Superfamily Hexapodoidea

Hexapodidae

Superfamily Leucosioidea

Iphiculidae

Leucosiidae

Superfamily Majoidea

Epialtidae

Hymenosomatidae

Inachidae

Majidae

Superfamily Palicoidea

Palicidae

Superfamily Parthenopoidea

Parthenopidae

Superfamily Pilumnoidea

Pilumnidae

Superfamily Portunoidea

Portunidae

Superfamily Retroplumoidea

Retroplumidae

Superfamily Trapezioidea

Trapeziidae

Superfamily Xanthoidea

Panopeidae

Xanthidae

Section Podotremata

Superfamily Cyclodorippoidea

Cyclodorippidae

Three species were identified using Tavares (1993). All are new for WA.

Krangalangia spinosa (Zarenkov, 1970)

MoV sp. 5024

Records: 5 specimens, 31°58'S, 848–1050 m

Distribution: N Australia; first record for S WA

Reference: Tavares (1993) [photo below]



Tymolus brucei Tavares, 1991

MoV sp. 5484

Records: 13 specimens, 21°00'S–35°04'S, 378–508 m

Distribution: N WA; first record for S WA

Reference: Tavares (1993)

Tymolus similis (Grant, 1905)

MoV sp. 5023

Records: many specimens, 22°04'S–35°22'S, 364–1050 m ✓

Distribution: SE Australia; first record for S WA

Reference: Tavares (1993) [photo below]



Cynomonidae

Of two species, one is a new record for Australia and the other a probable new species. Ahyong and Brown (2003) provided a key to Indo-West Pacific species.

Cynomonus andamanicus Alcock, 1905

MoV sp. 5025

Records: 1 specimen, 29°50'S, 408–427 m

Distribution: Indo-West Pacific; first record for Australia

Reference: Ahyong and Brown (2003)

Cynomonus sp. MoV 5001

Records: 12 specimens, 29°52'S–35°22'S, 401–1050 m

Distribution: new species

Reference: Ahyong and Brown (2003) [photo below]



Superfamily Dromioidea

Dromiidae

Three of the six species could be identified with the aid of McLay (1993). The others were placed in genera (one not previously recorded from Australia) using the same source but are not known species.

Austrodromidia insignis (Rathbun, 1923)

MoV sp. 3856

Records: 3 specimens, 27°48'S–35°16'S, 96–179 m

Distribution: S Australia

Reference: McLay (1993) [photo below]



Dromia wilsoni (Fulton & Grant, 1902)

MoV sp. 3854

Records: 16 specimens, 27°55'S–35°37'S, 95–253 m

Distribution: Indo-West Pacific and S Atlantic including Australia

Reference: McLay (1993) [photos below]




Epigodromia sp. MoV 5473

MoV sp. 5473

Records: 4 specimens, 35°20'S, 212–213 m

Distribution: new species

Reference: McLay (1993) [photo upper right]



Fultodromia nodipes (Guérin-Méneville, 1832)

MoV sp. 5029

Records: 3 specimens, 24°37'S–35°10'S, 97–100 m

Distribution: S Australia

Reference: McLay (1993) [photo below]



Fultodromia sp. MoV 5137

Records: 1 specimen, 27°48'S, 123–112 m

Distribution: new species

Reference: McLay (1993)

Takedromia sp. MoV 5003

Records: 5 specimens, 22°51'S–24°37'S, 100 m,

Distribution: new species, new record for genus in Australia

Reference: McLay (1993) [photo below]



Dynomenidae

One species was found (McLay, 1999), the first for southern WA.

Hirsutodynomena spinosa (Rathbun, 1911)

MoV sp. 5030

Records: 1 specimen, 27°03'S, 106 m

Distribution: Indo-West Pacific including N Australia; first record for S WA

Reference: McLay (1999) [photo below]



Superfamily Homoloidea

Homolidae

Among the seven species is a new Australian record and a possible new species (Guinot and Richer de Forges, 1995; Tan et al., 2000).

Dagnaudus petterdi (Grant, 1905)

MoV sp. 5038

Records: 9 specimens, 24°33'S–35°21'S, 364–490 m ✓

Distribution: E and S Australia, New Zealand, New Caledonia; new record for WA

Reference: Poore (2004) [photo below]



Homola orientalis Henderson, 1888

MoV sp. 5036

Records: 24 specimens, 21°59'S–31°37'S, 165–404 m ✓

Distribution: Indo-West Pacific including Australia

Reference: Poore (2004) [photo below]



Homologenus braueri Doflein, 1904

MoV sp. 5139

Records: 7 specimens, 21°55'S–31°43'S, 986–1295 m

Distribution: Indo-West Pacific including WA

Reference: Guinot and Richer de Forges (1995) [photo next page]



***Homologenus malayensis* Ihle, 1912**

MoV sp. 5039

Records: 24 specimens, 29°03'S–31°58'S, 848–1050 m

Distribution: West Pacific; first record for Australia

Reference: Guinot and Richer de Forges (1995) [photo below]



***Latreilopsis tetraspina* Dai & Chen, 1980**

MoV sp. 5035

Records: 1 specimen, 27°48'S, 123–112 m ✓

Distribution: WA

Reference: Guinot and Richer de Forges (1995)

***Paramolopsis boasi* Wood-Mason, 1891**

MoV sp. 5037

Records: 3 specimens, 21°00'S–22°04'S, 399–408 m, ✓

Distribution: Indo-West Pacific including Australia

Reference: Guinot and Richer de Forges (1995)

***Yaldwynopsis* sp. MoV 5004**

MoV sp. 5004

Records: 1 specimens, 31°37.27'S, 205–210 m ✓

Distribution: probable new species

Reference: Guinot and Richer de Forges (1995) [photo below]



Latreilliidae

Two species were identified confidently using Castro et al. (2003). One was recorded from Australia for the first time.

***Eplumula australiensis* (Henderson, 1888)**

MoV sp. 5040

Records: 11 specimens, 25°54'S–31°43'S, 100–253 m ✓

Distribution: Australia, New Zealand, New Caledonia

Reference: Williams (1982), Poore (2004) [photo below]



***Latreillia pennifera* Alcock, 1900**

MoV sp. 5041

Records: 25 specimens, 20°59'S–22°04'S, 100–408 m ✓

Distribution: Indo-West Pacific; first record for Australia

Reference: Castro et al. (2003) [photo below]



Superfamily Raninoidea

Raninidae

Of five species, four were well-known Australia species (Goeke, 1985; Dawson and Yaldwyn, 2000). The fifth was a species known previously from Japan-Philippines and now recorded from WA.

Cosmonotus grayi Adams & White, 1848

MoV sp. 5293

Records: 2 specimens, 20°59'S–27°48'S, 100–123 m

Distribution: Indo-West Pacific including NW Australia; first record for S WA

Reference: Sakai (1976: pl. 20, fig. 3) [photo below]



Lyreidus stenops Wood Mason, 1887

MoV sp. 5140

Records: 4 specimens, 21°58.41'S–22°4.28'S, 101–177 m ✓

Distribution: West Pacific; new record for Australia

Reference: Goeke (1985) [photo right]



Lyreidus tridentatus De Haan, 1841

MoV sp. 5295

Records: 17 specimens, 21°00'S–31°55'S, 201–414 m ✓

Distribution: Indo-West Pacific including Australia

Reference: Poore (2004) [photo below]



Notosceles serratifrons (Henderson, 1893)

MoV sp. 5294

Records: 21 specimens, 21°58'S–27°48'S, 106–166 m

Distribution: Indian Ocean including WA

Reference: Poore (2004) [photo next page]



Umalia trirufomaculata (Davie & Short, 1989)

MoV sp. 5296

Records: 12 specimens, 24°01'S–31°43'S, 100–123 m ✓

Distribution: N Australia; new record for S WA

Reference: Poore (2004) [photo below]



Section Eubrachyura

Subsection Heterotremata

Superfamily Aethroidea

Aethridae

Two species previously treated as members of Leucosiidae (Davie, 2002) were identified with reference to Alcock (1895), Miers (1876) and Griffin (1972). Both were found further south than hitherto known.

Actaeomorpha erosa Miers, 1877

MoV sp. 5061

Records: 1 specimen, 27°48'S, 98 m

Distribution: Indo-West Pacific including Qld and N WA; new record for S WA

Reference: Miers (1876), Alcock (1895) [photo below]



Drachiella sculpta (Haswell, 1879)

MoV sp. 5060

Records: 4 specimens, 20°59'S–21°59'S, 100–166 m

Distribution: N Australia; new record for S WA, here to greater depths than previously

Reference: Griffin (1972) [photo below]



Superfamily Calappoidea

Calappidae

Six species in two genera were found and identified using two papers by Galil (1993; 1997). One species could not be identified and another appeared a new Australian record. Three species are new for WA.

Calappa depressa Miers, 1886

MoV sp. 5016

Records: 7 specimens, 22°50'S–29°48'S, 95–114 m

Distribution: Indo-West Pacific including Australia

Reference: Galil (1997) [photos below]



Calappa lophos (Herbst, 1785)

MoV sp. 5017

Records: 3 specimens, 20°59'S–22°04'S, 100–107 m

Distribution: Indo-West Pacific including Australia

Reference: Galil (1997) [photos below]



Calappa pustulosa Alcock, 1896

MoV sp. 5018

Records: 7 specimens, 20°59'S–22°04'S, 100–177 m ✓

Distribution: West Pacific; new record for Australia

Reference: Galil (1997) [photos below]



Mursia australiensis Campbell, 1971

MoV sp. 5020

Records: 1 specimens, 31°36'S, 329–370 m

Distribution: E Australia; new record for WA

Reference: Galil (1993) [photos upper right]



Mursia microspina Davie & Short, 1989

MoV sp. 5019

Records: 3 specimens, 24°37'S–25°55'S, 100–120 m

Distribution: West Pacific including NE Australia; new record for WA

Reference: Galil (1993) [photos below]



Mursia sp. MoV 4988

Records: 6 specimens, 21°00'S–33°00'S, 387–428 m ✓

Distribution: new species similar to *M. musorstomi* Galil, 1993

Reference: Galil (1993) [photos below]



Superfamily Cancroidea

Atelecyclidae

Two species, both apparently undescribed were recorded. They could not be identified using Salva and Feldmann (2001).

Trichopeltarion sp. MoV 5135

Records: 40 specimens, 27°48'S–35°13'S, 364–494 m ✓

Distribution: new species different from those recorded from Tas. Seamounts

Reference: Poore et al. (1998) [photo below]



Trichopeltarion sp. MoV 5138

Mov sp. 5138

Records: 3 specimens, 28°59'S–35°04'S, 378–407 m

Distribution: new species similar to *T. wardi* Dell, 1968

Reference: Dell (1968) [photo below]



Superfamily Carpilioidea

Carpiliidae

A single species was recognised.

Carpilius convexus (Forskål, 1775)

MoV sp. 5080

Records: 1 specimen, 21°56'S, 134–132 m

Distribution: Indo-West Pacific including N Australia; new record for S WA

Reference: Serène (1984)

Superfamily Corystoidea

Corystidae

One of the two species (if correctly identified) is a new Australian record for a species previously described from Taiwan (Ng et al., 2000).

Gomeza bicornis Gray, 1831

MoV sp. 5022

Records: 1 specimen, 25°54'S, 100 m ✓

Distribution: Indo-West Pacific including S Australia

Reference: Hale (Hale, 1927)

Jonas cf. *choprai* Serène, 1971

MoV sp. 5021

Records: 3 specimens, 25°55'S–27°48'S, 123–112 m

Distribution: possible new species similar to *J. choprai* from Taiwan

Reference: Ng et al.(2000) [photo below]



Superfamily Dorippoidea

Dorippidae

All three species were identified to genus using Manning and Holthuis (1981).

Dorippe quadridens (Fabricius, 1793)

MoV sp. 5026

Records: 1 specimen, 22°04'S, 102 m

Distribution: Indo-West Pacific including Australia

Reference: Holthuis and Manning (1990)

Neodorippe callida Fabricius, 1798

MoV sp. 5027

Records: 1 specimen, 25°54'S, 100 m ✓

Distribution: Indo-West Pacific; new record for Australia (also known from Qld)

Reference: Holthuis and Manning (1990)

Paradorippe australiensis (Miers, 1884)

MoV sp. 5028

Records: 3 specimens, 24°37'S–25°54'S, 100–120 m ✓

Distribution: N Australia and W Papua; first record for S WA

Reference: Holthuis and Manning (1990),

Ethusidae

Four species were found but only one was tentatively identified using the keys and illustrations of Chen (1993). All are probable new species. The species of the family were previously treated as members of Dorippidae.

Ethusa cf. *granulosa* Ihle, 1916

MoV sp. 5006

Records: 2 specimens, 27°48'S–27°48'S, 112–436 m

Distribution: new species

Reference: Chen (1993: key)

Ethusa sp. MoV 5007

Records: 7 specimens, 21°56'S–31°43'S, (102) 848–1050 m

Distribution: new species

Reference: Chen (1993: key)

Ethusa sp. MoV 5008

Records: 1 specimen, 21°58'S, 726–732 m

Distribution: new species

Reference: Chen (1993: key)

Ethusina sp. MoV 5005

Records: 1 specimen, 31°58'S, 848–1050 m

Distribution: new species

Reference: Chen (1993: key) [photo below]



Superfamily Eriphioidea

Hypothalassiidae

The only records are probable juveniles of a large commercially exploited species (Koh and Ng, 2000). The genus was placed in Eriphiidae by Poore (2004) and other authors.

Hypothalassia acerba Koh & Ng, 2000

MoV sp. 5114

Records: 2 specimens, 31°37'S–35°22'S, 195–210 m

Distribution: S Australia

Reference: Koh and Ng (2000) [photo below]



Superfamily Goneplacoidea

Chasmocarcinidae

Two species of *Camatopsis* and one of *Megaesthius* were identified using Tesch (1918) and included in this family (rather than Goneplacidae) on the basis of arguments in Ng (1987).

Camatopsis rubida Alcock & Anderson, 1899

MoV sp. 5084

Records: 8 specimens, 21°58'S–22°04'S, 101–399 m

Distribution: Indo-West Pacific; new record for Australia

Reference: Sakai (1976)

Camatopsis sp. MoV 5086

Records: 1 specimen, 21°58'S, 373–382 m ✓

Distribution: new species

References: Sakai (1976), Tesch (1918)

Megaesthius sagaedae Rathbun, 1909

MoV sp. 5092

Records: 1 specimen, 22°04'S, 106–101 m

Distribution: Singapore; first record for Australia

Reference: Tesch (1918)

Euryplacidae

Both species are probable new species of *Heteroplax*. The genus has previously been included in Goneplacidae but we follow Ng et al. (2008) in placing them in Euryplacidae.

Heteroplax sp. MoV 4993

Records: 12 specimens, 22°04'S–27°55'S, 206–253 m ✓

Distribution: new species

References: Sakai (1976), Tesch (1918)

Heteroplax sp. MoV 4994

Records: many specimens, 21°59'S–22°02'S, 105–206 m ✓

Distribution: new species

References: Sakai (1976), Tesch (1918) [photo below]



Goneplacidae

The systematics of Goneplacidae and related families are difficult. Some species initially placed in this family were reassigned to other families (Chasmocarcinidae, Euryplacidae and Mathildellidae) on the basis of the arguments in Ng and Manuel-Santos (2007) and Ng (1987). Castro (2007) provided a key to genera of Goneplacinae, a subfamily used as a family here, but not all species could be identified confidently to genus or species.

Carcinoplax sp. MoV 4996

Records: 1 specimen, 21°59'S, 166 m

Distribution: new species

Reference: Guinot (1989)

Carcinoplax sp. MoV 4998

Records: 6 specimens, 27°48'S–35°22'S, 416–695 m

Distribution: new species

Reference: Guinot (1989)

Notonyx nitidus Milne Edwards, 1873

MoV sp. 5088

Records: 2 specimens, 22°51'S–22°02'S, 100–105 m

Distribution: West Pacific; first record for Australia

Reference: Clark and Ng (2006) [photo below]



Psopheticus stridulans Wood-Mason, 1892

MoV sp. 5032

Records: 23 specimens, 21°58'S–33°00'S, 373–423 m, ✓

Distribution: Indo-West Pacific; first record for Australia

Reference: Sakai (1976) [photo below]



Pycnoplax bispinosa (Rathbun, 1914)

MoV sp. 4991

Records: 18 specimens, 21°58'S–22°04'S, 170–206 m ✓

Distribution: first record for Australia

References: Guinot (1989), Castro (2007)

Pycnoplax meridionalis (Rathbun, 1923)

MoV sp. 3862

Records: 7 specimens, 31°37'S–35°23'S, 147–776 m

Distribution: S Australia

References: Poore (2004) as *Carcinoplax meridionalis*, Castro (2007) [photo below]



Pycnoplax victoriensis (Rathbun, 1923)

MoV sp. 5031

Records: 3 specimens, 28°59'S–35°21'S, 389–704 m

Distribution: SE Australia; first record for S WA

Reference: Poore (2004) as *Carcinoplax victoriensis*, Castro (2007)

Pycnoplax cf. *surugensis* (Rathbun, 1932)

MoV sp. 4992

Records: 14 specimens, 21°55'S–31°57'S, 848–1295 m

Distribution: probable new species close to Japanese species

Reference: Guinot (1989), Castro (2007) [photo below]



Pycnoplax sp. MoV 5124

Records: 1 specimen, 21°59'S, 166 m

Distribution: new species

Reference: Guinot (1989)

Mathildellidae

Two species previously considered members of Goneplacidae were found. Family placement follows Ng et al. (2008).

Mathildella serrata (Sakai, 1974)

MoV sp. 5112

Records: 13 specimens, 27°55'S–35°22'S, 205–915 m ✓

Distribution: West Pacific; new record for Australia, also known from SE Australia

Reference: Ng and Chan (2000) [photo below]



Platypilumnus soelae Garth, 1987

MoV sp. 5033

Records: 2 specimens, 21°58'S, 356–324 m

Distribution: N WA; first record for S WA

Reference: drawing from Garth (1987)

Mathildellid sp. MoV 4997

Records: 1 specimen, 31°58'S, 848–1050 m

Distribution: juvenile of new species difficult to place in genus; may belong in Goneplacidae

Reference: Tesch (1918) [photo below]



Superfamily Hexapodidae

Hexapodidae

The only species has been recorded before only from Japan to Indonesia (Manning and Holthuis, 1981).

Hexaplax megalops Dolflein, 1904

MoV sp. 5034

Records: 11 specimens, 21°00'S–22°04'S, 387–408 m ✓

Distribution: West Pacific; new record for Australia

References: Manning and Holthuis (1981: key), Sakai (1976) [photo below]



Superfamily Leucosioidea

Iphiculidae

The single species was previously treated as a member of Leucosiidae. It is a new record for Australia.

Iphiculus spongiosus Adams & White, 1848

MoV sp. 5113

Records: 10 specimens, 20°59'S–22°04'S, 100–107 m ✓

Distribution: Indo-West Pacific; new record for Australia

Reference: Chen (1989)

Leucosiidae

Numerous species are known from Australia but not all in these samples could be identified to species. The Western Australian fauna was reviewed by Tyndale-Biscoe and George (1962). Tentative identifications were made using names of species described from more northern parts of the West Pacific.

The family was represented by 30 species (many in just one sample) in ten genera. For some genera the literature is scattered but Alcock (1895) and Sakai (1976) are useful to identify genera. Chen (1989), Tan and Ng (1996) and Tan (1996) included similar or the same species. Thirteen species are new (42%) and four are new records for Australia.

Arcania cornuta (MacGilchrist, 1905)

MoV sp. 5045

Records: 4 specimens, 21°58'S–25°55'S, 120–177 m

Distribution: Indo-West Pacific including Qld; new record for WA

References: Chen (1989), Galil (2001a) [photo below]



Arcania elongata Yokoya, 1933

MoV sp. 5042

Records: 3 specimens, 21°59'S–30°59'S, 100–166 m

Distribution: Japan, Qld. NSW; first record for WA

Reference: Galil (2001a) [photo below]



Arcania gracilis (Henderson, 1893)

MoV sp. 5047

Records: 4 specimens, 20°59'S–21°59'S, 100–166 m

Distribution: Indo-West Pacific including WA; first record for S WA

Reference: Galil (2001a)



Arcania muricata Galil, 2001

MoV sp. 5046

Records: 4 specimens, 20°59'S–21°58'S, 100–177 m

Distribution: Indo-West Pacific including NT; first record for S WA

Reference: Galil (2001a) [photo below]



Arcania sp. MoV 4980

Records: 3 specimens, 27°48'S–29°48'S, 96–114 m

Distribution: new species like *A. sagmiensis* from Japan

Reference: Galil (2001a) [photo below]



Arcania novemspinosa (Adams & White, 1849)

MoV sp. 5043

Records: 1 specimen, 25°55'S, 120 m

Distribution: Indo-West Pacific including WA

Reference: Galil (2001a) [photo below]



Arcania sp. MoV 4987

Records: 2 specimens, 22°50'S–24°37'S, 100 m

Distribution: new species like *A. septemspinosa*

Reference: Galil (2001a) [photo below]



Arcania septemspinosa (Fabricius, 1787)

MoV sp. 5044



Ebalia tuberculosa (Milne Edwards, 1873)

MoV sp. 0710

Records: many specimens, 21°59'S–35°22'S, 212–539 m ✓

Distribution: Indo-West Pacific including Australia

Reference: Poore (2004) [photo next page]




Ebalia sp. MoV 4981

Records: 6 specimens, 22°50'S–24°01'S, 100 m

Distribution: new species like *E. dimorphoides*

Reference: Chen (1989) [photo below]



Ebalia sp. MoV 4989

Records: 3 specimens, 20°59'S–21°58'S, 100–107 m

Distribution: new species

Reference: Chen (1989) [photo below]



Ebalia sp. MoV 4990

Records: 1 specimen, 28°59'S, 180–183 m

Distribution: new species

Reference: Chen (1989) [photo below]



Leucosia haematosticta Adams & White, 1849

MoV sp. 5053

Records: 1 specimen, 25°54'S, 100 m

Distribution: Indo-West Pacific including Australia; new record for S WA

Reference: Poore (2004) [photo below]



Leucosia ocellata Bell, 1855

MoV sp. 5064

Records: 1 specimen, 21°58'S, 177–170 m

Distribution: N Australia; new record for S WA

Reference: Campbell and Stephenson (1970) [photo below]



Leucosia whitei Bell, 1855

MoV sp. 5052

Records: 1 specimen, 20°59'S, 100 m

Distribution: Indo-West Pacific including Australia

Reference: Arnold and George (1987) [photo below]



***Leucosia* sp. MoV 4985**

Records: 1 specimen, 22°04'S, 102 m

Distribution: new species like *L. foresti*

Reference: Chen (1989) [photo below]



Merocryptus lambriiformis Milne Edwards, 1873

MoV sp. 3864

Records: 6 specimens, 35°20'S–35°22'S, 161–213 m

Distribution: West Pacific including Australia

Reference: Poore (2004) [photo below]



Myra curtimana Galil, 2001

MoV sp. 5050

Records: 15 specimens, 21°57'S–35°25'S, 100–1031 m ✓

Distribution: West Pacific including WA

Reference: Galil (2001b) [photo below]



***Myra* sp. MoV 4982**

Records: 1 specimens, 27°48'S, 123–112 m

Distribution: new species

Reference: Galil (2001b) [photo below]



***Myra* sp. MoV 4983**

Records: 3 specimens, 20°59'S, 100 m

Distribution: new species

Reference: Galil (2001b) [photo below]



Myrine kessleri (Paulson, 1875)

MoV sp. 5051

Records: 2 specimens, 20°59'S–21°57'S, 100–114 m

Distribution: Indo-West Pacific including N Australia; first record for S WA

Reference: Galil (2001b) [photo below]



Oreophorus reticulatus Adams & White, 1849

MoV sp. 5062

Records: 2 specimens, 20°59'S–27°48'S, 100–123 m ✓

Distribution: Indo-West Pacific; new record for Australia

Reference: Tan and Ng (1996) [photo below]



Parilia major Sakai, 1961

MoV sp. 5055

Records: 1 specimen, 22°04'S, 396–391 m

Distribution: Japan; new record for Australia (doubtful identification)

Reference: Sakai (1976) [photo below]



Philyra sp. MoV 4984

Records: 1 specimen, 24°01'S, 100 m

Distribution: new species

Reference: Poore (2004) [photo below]



Randallia eburnea Alcock, 1896

MoV sp. 5048

Records: 16 specimens, 21°59'S–35°21'S, 100–404 m

Distribution: Indo-West Pacific including Australia

Reference: Chen (1989) [photo below]



Randallia pustuloides Sakai, 1961

MoV sp. 5049

Records: 2 specimens, 21°58'S, 373–382 m

Distribution: Japan, Philippines; new record for Australia

Reference: Chen (1989) [photo below]



Randallia sp. MoV 4977

Records: 1 specimen, 22°37'S, 355–382 m

Distribution: new species

Reference: Chen (1989) [photo next page]



***Randallia* sp. MoV 4978**

Records: 40 specimens, 20°59'S–27°48'S, 100–166 m ✓

Distribution: new species

Reference: Chen (1989) [photo below]



***Randallia* sp. MoV 4979**

Records: 1 specimen, 21°58'S, 107 m

Distribution: new species

Reference: Chen (1989)

***Randallia* sp. MoV 4986**

Records: 1 specimen, 21°59'S, 166 m

Distribution: new species similar to *R. speciosa*

Reference: Chen (1989)

Superfamily Majoidea

Epialtidae

The family name Epialtidae is used to include what were previously treated as subfamilies Epialtinae and Pisinae of Majidae. We follow the arrangement of Ng et al. (2008). Taxonomy follows Griffin and Tranter (1986) who reviewed the fauna and provided keys to Majidae in the broadest sense. Reference to older and more recent papers was required for some genera (Griffin, 1970, 1973; Guinot and Richer de Forges, 1982, 1985). The 14 species include several new records, one from Australia, and three new species under study by B. Richer de Forges.

***Austrolibinia gracilipes* (Miers, 1879)**

MoV sp. 5162

Records: 4 specimens, 20°59'S, 100 m

Distribution: Indonesia, PNG, N Australia; first record for S WA

Reference: Miers (1879: pl. 4, fig. 4) [photo below]



***Griffinia lappacea* (Rathbun, 1918)**

MoV sp. 5173

Records: 1 specimen, 34°00'S, 467–490 m

Distribution: Australia

Reference: Griffin and Tranter (1986) [photo below]



***Hyastenus convexus* Miers, 1884**

MoV sp. 5169

Records: 24 specimens, 20°59'S–28°58'S, 95–120 m ✓

Distribution: Indo-West Pacific, N Australia; first record for S WA

Reference: Griffin and Tranter (1986) [photo below]



***Lahaina agassizii* (Rathbun, 1902)**

MoV sp. 5172

Records: 18 specimens, 22°50'S–33°58'S, 96–100 m ✓

Distribution: Indo-West Pacific including Australia

Reference: Griffin and Tranter (1986) [photo below]



***Naxioides robillardi* (Miers, 1882)**

MoV sp. 5174

Records: 1 specimen, 21°58'S, 177–170 m

Distribution: Indo-West Pacific including E Australia; first record for WA

Reference: Poore (2004) [photo below]



***Naxioides taurus* (Pocock, 1890)**

MoV sp. 5165

Records: 1 specimen, 21°01'S, 93 m

Distribution: Indo-West Pacific including N Australia; first record for WA

Reference: Griffin and Tranter (1986) [photo upper right]



***Naxioides tenuirostris* (Haswell, 1880)**

MoV sp. 5164

Records: 1 specimen, 27°55'S, 253 m

Distribution: Indo-West Pacific including N Australia; first record for WA

Reference: Griffin and Tranter (1986) [photo below]




***Phalangipus filiformis* Rathbun, 1916**

MoV sp. 5160

Records: 2 specimens, 20°59'S–22°4'S, 100 m

Distribution: Indo-West Pacific including N Australia; first record for S WA

Reference: Griffin (1973) [photo below]



Phalangipus hystrix (Miers, 1884)

MoV sp. 5161

Records: 27 specimens, 21°58'S–27°48'S, 100–166 m ✓

Distribution: Indo-West Pacific including WA

Reference: Griffin (1973) [photos below]




Rochinia fultoni (Grant, 1905)

MoV sp. 3895

Records: 1 specimen, 27°55'S, 253 m

Distribution: SE Australia; first record for WA

Reference: Poore (2004) [photo below]



Rochinia* aff. *luzonica (Rathbun, 1916)

MoV sp. 5168

Records: 5 specimens, 29°00'S–31°37'S, 329–439 m

Distribution: new species

Reference: Griffin (1976) [photo below]



Rochinia strangeri Serène & Lohavanijaya, 1973

MoV sp. 5538

Records: 1 specimens, 29°3.39'S, 1000–1037 m

Distribution: S China Sea; first record for Australia

Reference: Serène and Lohavanijay (1973) (det. B. Richer de Forges)

***Rochinia* sp. MoV 5119**

Records: numerous specimens, 29°52'S–35°04'S, 329–414 m

Distribution: new species

Reference: Griffin and Tranter (1986)

***Rochinia* sp. MoV 5136**

Records: 3 specimens, 21°58'S–23°59'S, 324–411 m

Distribution: new species close to "*Sphenocarcinus carbunculus* Rathbun, 1906" from Hawaii

Reference: Rathbun (1906)

***Epialtid* sp. MoV 5134**

Records: 1 specimen, 33°58'S, 96 m

Distribution: new species, genus uncertain, possibly *Thycaophrys*

Reference: Griffin and Tranter (1986) [photo below]



Hymenosomatidae

One of the two species could not be identified beyond genus from Lucas (1980) or Ng and Chuang (1996).

Halicarcinus sp. MoV 5002

Records: 3 specimens, 28°58'S–35°10'S, 86–107 m

Distribution: new species

References: Lucas (1980), Ng and Chuang (1996) [photo below]



Trigonoplax longirostris McCulloch, 1908

MoV sp. 1678

Records: 1 specimen, 31°43'S, 102 m

Distribution: Australia; recorded at depth

Reference: Lucas (1980) [photo below]



Inachidae

The Inachidae were treated as a subfamily of Majidae in earlier literature but are elevated to family rank in this report as advocated by Ng et al. (2008). Taxonomy follows Griffin and Tranter (1986) who reviewed the fauna and provided keys to Majidae in the broadest sense. Inachidae include 20 species of which two are new and three are new Australian records. Reference to older and more recent papers was required for some genera (Griffin, 1970, 1973).


Achaeus brevirostris (Haswell, 1879)

MoV sp. 5347

Records: 3 specimens, 21°58'S–27°48'S, 100–123 m

Distribution: Indo-West Pacific including Australia

Reference: Griffin and Tranter (1986) [photo below]




Achaeus curvirostris (Milne Edwards, 1873)

MoV sp. 3851

Records: 4 specimens, 27°48'S–33°02'S, 95–123 m

Distribution: Indo-West Pacific including Australia

Reference: Griffin and Tranter (1986) [photo below]



Achaeus lacertosus Stimpson, 1857

MoV sp. 5150

Records: 4 specimens, 21°58'S–27°48'S, 100–123 m

Distribution: Indo-West Pacific including Australia; new record for S WA

Reference: Griffin and Tranter (1986)

Achaeus sp. MoV 5122

Records: 2 specimens, 22°50'S–27°48'S, 123–100 m

Distribution: new species

Reference: Griffin and Tranter (1986)

***Achaeus* sp. MoV 5123**

Records: 3 specimens, 27°48.29'S, 123–112 m ✓

Distribution: recorded as undescribed species by Griffin

Reference: Griffin (1970)


***Camposcia retusa* Latreille, 1829**

MoV sp. 5151

Records: 1 specimens, 31°43.28'S, 102 m

Distribution: Indo-West Pacific including Australia; new record for S WA

Reference: Griffin and Tranter (1986) [photo below]



***Dorhynchus ramusculus* (Baker, 1906)**

MoV sp. 5159

Records: 10 specimens, 29°52.04'S–35°21.53'S, 212–490 m

Distribution: New Zealand, S Australia

References: Poore (2004) [photo below]



***Cyrtomaia maccullochi* Rathbun, 1918**

MoV sp. 5146

Records: 34 specimens, 27°08'S–35°12'S, 378–728 m

Distribution: Indo-West Pacific including Australia

Reference: Griffin and Tranter (1986) [photo below]



***Dumea latipes* (Haswell, 1880)**

MoV sp. 1338

Records: 3 specimens, 31°43'S, 102 m

Distribution: S Australia

References: Poore (2004)

***Ephippias endeavouri* Rathbun, 1918**

MoV sp. 5158

Records: 4 specimens, 31°43'S–35°22'S, 102–196 m

Distribution: SE and SW Australia

References: Poore (2004) [photo below]



***Cyrtomaia murrayi* Miers, 1886**

MoV sp. 5147

Records: 3 specimens, 27°55'S–31°37'S, 252–404 m

Distribution: Indo-West Pacific including Australia

Reference: Griffin and Tranter (1986) [photo upper right]


Grypachaeus hyalinus Alcock & Anderson, 1894

MoV sp. 5148

Records: 2 specimens, 20°59.05'S–24°01'S, 100 m

Distribution: Indian Ocean, new Australian record

References: Griffin and Tranter (1986) [photo below]



Oncinopus aranea De Haan, 1839

MoV sp. 5154

Records: 1 specimen, 27°08'S, 414–405 m

Distribution: W Pacific, including Australia

References: Poore (2004) [photo below]



Oncinopus* cf. *angustifrons Takeda & Miyake, 1969

MoV sp. 5120

Records: 3 specimens, 24°01'S–31°43'S, 100–183 m

Distribution: Japan, Phillipines, new Australian record if correctly identified

References: Griffin and Tranter (1986) [photo below]



Oncinopus neptunus Adams & White, 1848

MoV sp. 5155

Records: 6 specimens, 24°37'S–, 31°43'S, 95–100 m

Distribution: Indo-West Pacific, including Australia

References: Poore (2004) [photo upper right]

Physachaeus ctenurus Alcock, 1895

MoV sp. 5149

Records: 56 specimens, 29°52'S–35°21'S, 364–528 m

Distribution: Andaman Sea, new Australian record

References: Griffin and Tranter (1986) [photos below]



Platymaia wyvillethomsoni Miers, 1886

MoV sp. 5157

Records: 24 specimens, 23°59'S–35°12'S, 364–431m

Distribution: West Pacific, Australia

References: Guinot and Richer de Forges (1985) [photo below]



***Platymaia fimbriata* Rathbun, 1916**

MoV sp. 5156

Records: 1 specimen, 21°58'S, 726–732 m

Distribution: West Pacific, Australia

References: Guinot and Richer de Forges (1985)

***Sunipea indicus* (Alcock, 1895)**

MoV sp. 5171

Records: 14 specimens, 22°51'S–29°48'S, 85–123m

Distribution: Andaman Sea, new Australian record

References: Griffin and Tranter (1986) [photo below]



Majidae

The family name Majidae is used in this report in the narrow sense advocated by Ng et al. (2008). The subfamilies used by, for example Davie (2002), are treated as families: Epialtinae and Pisinae together as Epialtidae; Planoterginae and Majinae as Majidae; and Inachinae as Inachidae. Taxonomy follows Griffin and Tranter (1986) who reviewed the fauna and provided keys to Majidae in the broadest sense. Forty-eight species of majids in the broadest sense were recognised of which 14 belong to Majidae s. s. Five species are new records of Indo-West Pacific species in Australia and four are new species.


***Entomonyx depressus* Sakai, 1974**

MoV sp. 5167

Records: 5 specimens, 22°50'S, 100 m

Distribution: Japan; new record for Australia

Reference: Griffin and Tranter (1986) [photo below]



***Entomonyx spinosus* Miers, 1884**

MoV sp. 5166

Records: 13 specimens, 22°50'S–35°21'S, 100–179 ✓

Distribution: Indo-West Pacific including N WA; first record for S WA

Reference: Griffin and Tranter (1986) [photo below]




***Leptomithrax globifer* Rathbun, 1918**

MoV sp. 5144

Records: 17 specimens, 35°37'S–35°22'S, 99–196 m

Distribution: S Australia; first positive record for S WA

Reference: Poore (2004) [photo below]




***Leptomithrax* sp. MoV 5121**

Records: 3 specimens, 28°59'S–34°49'S, 50–232 m

Distribution: new species

Reference: Griffin and Tranter (1986) [photo below]



***Leptomithrax sternocostulatus* (Milne Edwards, 1851)**

MoV sp. 0703

Records: 5 specimens, 28°58'S–35°37'S, 86–106 m ✓

Distribution: S Australia

Reference: Poore (2004) [photo below]



***Maja confragosa* Griffin & Tranter, 1986**

MoV sp. 5152

Records: 1 specimen, 22°37'S, 355–382 m

Distribution: Indonesia; new record for Australia

Reference: Griffin and Tranter (1986) [photo below]



***Leptomithrax* sp. MoV 5133**

Records: 1 specimen, 27°48'S, 123–112 m

Distribution: new species

Reference: Griffin and Tranter (1986) [photo upper right]

Maja gibba Alcock, 1895

MoV sp. 5145

Records: 1 specimen, 22°37'S, 355–382 m

Distribution: West Pacific; first record for Australia

Reference: Griffin and Tranter (1986) [photo below]



Maja suluensis Rathbun, 1916

MoV sp. 5143

Records: 1 specimen, 24°02'S, 100 m

Distribution: Indonesia; first record for Australia

Reference: Griffin and Tranter (1986) [photo below]




Planotergum mirabile Balss, 1935

MoV sp. 5153

Records: 1 specimen, 28°58'S, 85 m ✓

Distribution: Indo-West Pacific including Australia

Reference: Davie (2002) [photo upper right]



Pristatopus brevispinosus Yokoya, 1933

MoV sp. 5298

Records: 5 specimens, 31°36'S, 329–370 m

Distribution: Japan; first record for Australia

Reference: Sakai (1976: 251) [photo below]



Pristatopus occidentalis Griffin, 1970

MoV sp. 5163

Records: 12 specimens, 27°55'S–35°20'S, 179–253 m

Distribution: WA endemic

Reference: Poore (2004) [photo below]



***Prismatopus* sp. MoV 5125**

Records: 1 specimen, 24°37'S, 100 m ✓

Distribution: new species

Reference: Griffin and Tranter (1986) [photo below]



Superfamily Palicoidea

Palicidae

In spite of the recent thorough study by Castro (2000), two of the five species could not be confidently identified beyond genus.

***Micropalicus vietnamensis* (Zarenkov, 1968)**

MoV sp. 5058

Records: 5 specimens, 21°58'S–22°00'S, 170–754 m

Distribution: West Pacific, NW Australia

References: Castro and Davie (2003) [photo below]



***Neopalicus jukesii* (White, 1847)**

MoV sp. 5000

Records: 1 specimen, 27°48'S, 123–112 m

Distribution: West Pacific, NW Australia

References: Castro and Davie (2003)

***Paliculus kyusyuensis* (Yokoya, 1933)**

MoV sp. 5057

Records: 5 specimens, 23°59'S–24°33'S, 388–412 m

Distribution: Indo-West Pacific, Qld, Australia


References: Castro and Davie (2003)

***Parapalicus* sp. MoV 4999**

Records: 25 specimens, 20°59'S–22°04'S, 100–177 m

Distribution: new species

References: Castro (2000) [photo below]



Pseudopalicus macromeles Castro, 2000

MoV sp. 5056

Records: 10 specimens, 27°55'S–35°20'S, 194–252 m ✓

Distribution: Australia

References: Castro (2000) [photo below]



Superfamily Parthenopoidea

Parthenopidae

Nine species were collected, of which only three could be confidently identified. Generic placement was made with reference to Sakai (1976) whose keys reflect those in Flipse (1930). Species identifications referred to Ng (1996), Garth and Davie (1995), Davie and Turner (1994) and Ahyong (2008).

Aulacolambrus sp. MoV 5014

Records: 2 specimens, 20°59'S, 100 m

Distribution: new species like *A. sibogae*

Reference: Flipse (1930)

Garthambrus cf. *lacunosa* (Rathbun, 1906)

MoV sp. 5011

Records: 1 specimen, 31°36'S, 329–370 m

Distribution: new species close to Hawaiian *G. lacunosa* and *G. tani* Ahyong, 2008

References: Ng (1996), Ahyong (2008)

Garthambrus cf. *stellatus* (Rathbun, 1906)

MoV sp. 5063

Records: 1 specimen, 22°37'S, 355–382 m

Distribution: identification of Hawaiian species uncertain

Reference: Ng (1996) [photo below]



Parthenope chondrodes Davie & Turner, 1994

MoV sp. 5010

Records: 7 specimens, 20°59'S–25°55'S, 100–120 m

Distribution: WA; new record for S WA

Reference: Davie and Turner (1994) [photo below]



***Platylambrus validus* De Haan, 1837**

MoV sp. 5065

Records: 2 specimens, 21°59'S–31°55'S, 165–232 m

Distribution: West Pacific including NE Australia; first record for WA

Reference: Campbell and Stephenson (1970)

***Pseudolambrus* sp. MoV 5009**

Records: 18 specimens, 22°50'S–27°48'S, 123–100 m ✓

Distribution: new species like *P. beaumonti*

Reference: Sakai (1976: 276, key)

***Rhinolambrus* sp. MoV 5012**

Records: 1 specimen, 27°48'S, 123–112 m ✓

Distribution: new species like *R. spinifer*

Reference: Flipse (1930)

***Thyrolambrus excavatus* Baker, 1905**

MoV sp. 5064

Records: 8 specimens, 27°48'S–35°10'S, 85–169 m

Distribution: S Australia; new record for WA

Reference: Baker (1905) [photo below]



***Parthenopid* sp. MoV 5015**

Records: 2 specimens, 35°11'S, 157–147 m

Distribution: new species not readily assigned to genus

Reference: Flipse (1930)

Superfamily Pilumnoidea

Pilumnidae

Three subfamilies were represented by 21 species. Half (ten species) are probably new species. Genera are listed alphabetically and subfamily names appear after species names.

Subfamily Eumedoninae was represented by one well-known species.

Subfamily Pilumninae included 15 species of which seven could be provisionally identified by P. Davie. Five are new records for WA. Where no reference is given, identification relies on P. Davie's unpublished notes.

Subfamily Rhizopinae was represented by five species, two identifiable to species but neither previously recorded from Australia (Tesch, 1918; Ng, 1987).

***Bathypilumnus pugilator* (Milne Edwards, 1873)**
(Pilumninae)

MoV sp. 5095

Records: 2 specimens, 24°01'S, 100 m

Distribution: New Caledonia, Qld; first record for WA

Reference: Davie (1989)

***Caecopilumnus piroculatus* (Rathbun, 1911)**
(Rhizopinae)

MoV sp. 5090

Records: 1 specimen, 27°48'S, 98 m

Distribution: Indonesia; first record for Australia

***Cryptolutea arafurensis* Davie & Humpherys, 1997**
(Rhizopinae)

MoV sp. 5085

Records: 1 specimen, 22°04'S, 106–101 m

Distribution: N Australia; first record for S WA

Reference: Davie and Humpherys (1997)

***Eumedonus niger* Milne Edwards, 1834 (Eumedoninae)**

MoV sp. 5111

Records: 2 specimens, 27°03'S–27°48'S, 106–123 m

Distribution: West Pacific including Australia

Reference: Chia and Ng (2000) [photo below]



***Heteropilumnus* sp. MoV 5101 (Rhizopinae)**

Records: 3 specimens, 25°54'S–35°22'S, 100–196 m ✓

Distribution: new species

Reference: Ng (1987)

***Lophoplax* sp. MoV 5105 (Pilumninae)**

Records: 1 specimen, 21°01'S, 93 m

Distribution: synonymous with a new genus and species from N Australia (P. Davie, pers. comm.)

Reference: Tesch (1918)

***Mertonia lanka* Laurie, 1906 (Rhizopinae)**

MoV sp. 5091

Records: 1 specimen, 24°01'S, 100 m

Distribution: Indian Ocean; first record for Australia

***Paraselwynia* sp. MoV 5089 (Rhizopinae)**

Records: 1 specimen, 33°58'S, 96 m

Distribution: new species, generic assignment problematic

Reference: Tesch (1918)

***Pilumnopeus* sp. MoV 5106 (Pilumninae)**

Records: 1 specimen, 22°02'S, 106 m

Distribution: new species

***Pilumnus* cf. *haswelli* De Man, 1888 (Pilumninae)**

MoV sp. 5104

Records: 1 specimen, 21°01'S, 93 m

Distribution: first record for Australia if correctly identified

***Pilumnus* cf. *hirsutus* Stimpson, 1858 (Pilumninae)**

MoV sp. 5098

Records: 4 specimens, 20°59'S–27°03'S, 100–414 m

Distribution: new record for Australia

***Pilumnus kingstoni* (Rathbun, 1923) (Pilumninae)**

MoV sp. 5097

Records: 8 specimens, 27°55'S–35°22'S, 105–253 m ✓

Distribution: S Australia; first record for WA

Reference: Poore (2004) [photo below]



***Pilumnus* cf. *propinquus* Nobili, 1905 (Pilumninae)**

MoV sp. 5297

Records: 2 specimens, 25°54'S, 100 m

Distribution: new record for Australia

***Pilumnus* cf. *schellenbergi* Balss, 1933 (Pilumninae)**

MoV sp. 5100

Records: 2 specimens, 21°59'S, 166 m

Distribution: new record for Australia [photo below]



***Pilumnus* cf. *spinicarpus* Grant & McCulloch, 1906 (Pilumninae)**

Records: 24 specimens, 21°57'S–29°48'S, 100–183 m ✓

Distribution: N Australia; new record for S WA if correctly identified [photo below]



***Pilumnus* sp. MoV 5094 (Pilumninae)**

Records: 1 specimens, 21°59'S, 166 m

Distribution: new species

***Pilumnus* sp. MoV 5099 (Pilumninae)**

MoV sp. 5099

Records: 4 specimens, 20°59'S–28°59'S, 100–183 m ✓

Distribution: new species [photo below]



***Pilumnus* sp. MoV 5103 (Pilumninae)**

Records: 1 specimen, 21°59'S, 166 m

Distribution: new species

***Pilumnus* sp. MoV 5474 (Pilumninae)**

Records: 1 specimen, 24°37'S, 100 m

Distribution: new species

Pilumnus sp. MoV 5475 (Pilumninae)

Records: 1 specimen, 27°48'S, 123–112 m

Distribution: new species

Pilumnid sp. MoV 4995 (Pilumninae)

Records: 1 specimen, 31°58'S, 848–1050 m

Distribution: new species [photo below]



Superfamily Portunoidea

Portunidae

Twenty-five species of swimming crabs were found, four not identifiable to species and one probably belonging to a new genus according to V. Spiridonov who examined some specimens. Half of the species are widespread in the Indo-West Pacific but only a few are newly recorded from Australia or WA. Identification was largely possible with reference to Stephenson (1972) and the earlier papers by this author but Wee and Ng (1995) was useful for *Charybdis* in particular and Davie and Crosnier (2006) for a recently described species.

Charybdis (Charybdis) miles (De Haan, 1835)

MoV sp. 5127

Records: 4 specimens, 21°58'S–21°59'S, 165–177 m

Distribution: Indo-West Pacific including N and E Australia; first record for WA

Reference: Poore (2004) [photo below]



Charybdis (Goniohellenus) hongkongensis Shen, 1934

MoV sp. 5190

Records: 1 specimen, 24°37'S, 100 m

Distribution: West Pacific; first record for Australia

Reference: Wee and Ng (1995) [photo below]



Echinolatus poorei Davie & Crosnier, 2006

MoV sp. 5141

Records: 22 specimens, 34°53'S–35°22'S, 95–484 m

Distribution: S Australia; first record for WA

Reference: Davie and Crosnier (2006) [photo below]



Libystes paucidentatus Stephenson & Campbell, 1960

MoV sp. 5188

Records: 1 specimen, 21°58'S, 177–170 m

Distribution: New Guinea, Qld; first record for WA

Reference: Stephenson (1972) [photo below]



Liocarcinus corrugatus (Pennant, 1777)

MoV sp. 5128

Records: 10 specimens, 24°37'S–27°48'S, 96–123 m

Distribution: Indo-West Pacific including Australia

Reference: Poore (2004)



Lissocarcinus orbicularis Dana, 1852

MoV sp. 5441

Records: 1 specimen, 22°50'S, 100 m

Distribution: Indo-West Pacific including Australia (from gut of holothurian)

Reference: Sakai (1976) [photo upper right]



Lupocyclus philippinensis Semper, 1880

MoV sp. 5130

Records: 14 specimens, 20°59'S–24°37'S, 100–107 m ✓

Distribution: Indo-West Pacific including NE Australia; first record for WA

Reference: Leene (1940) [photo below]



Lupocyclus quinquedentatus Rathbun, 1906

MoV sp. 5142

Records: 1 specimen, 25°54'S, 100 m

Distribution: West Pacific; first record for Australia

Reference: Leene (1940) [photo below]



***Lupocyclus* sp. aff. *tugelae* Barnard, 1950**

MoV sp. 5185

Records: 6 specimens, 21°59'S–27°48'S, 100–166 m

Distribution: new species close to *L. tugelae* (Indo-West Pacific including N WA)

Reference: Barnard (1950) [photo below]



***Nectocarcinus spinifrons* Stephenson, 1961**

MoV sp. 5129

Records: 12 specimens, 24°37'S–33°02'S, 95–102 m

Distribution: SW Australia

Reference: Poore (2004) [photo below]



***Ovalipes elongatus* Stephenson & Rees, 1968**

MoV sp. 5192

Records: 1 specimen, 35°21'S, 91 m

Distribution: New Zealand, Lord Howe; first record for WA

Reference: Stephenson (1972) [photo upper right]



***Ovalipes iridescens* (Miers, 1886)**

MoV sp. 5132

Records: 79 specimens, 22°37'S–27°48'S, 355–1010 m

Distribution: Indo-West Pacific including Vic.; first record for WA

Reference: Stephenson (1972) [photo below]



***Parathranites orientalis* (Miers, 1886)**

MoV sp. 5031

Records: many specimens, 21°59'S–30°59'S, 100–183 m ✓

Distribution: Indo-West Pacific including E Australia; first record for WA

Reference: Stephenson (1972) [photo next page]



Parathranites sp. MoV 5290

Records: 7 specimens, 22°50'S, 100 m

Distribution: new species? (det. V. Spiridonov) [photo below]



Portunus aff. *argentatus* (Milne Edwards, 1861)

MoV sp. 5287

Records: 45 specimens, 21°57'S–22°04'S, 101–107 m

Distribution: new species? (det. V. Spiridonov) [photo below]



Portunus (Monomia) haanii (Stimpson, 1858)

MoV sp. 5125

Records: 18 specimens, 22°04'S–33°58'S, 96–102 (1085) m

Distribution: Indo-West Pacific including Australia

Reference: Poore (2004) [photo below]



Portunus (Xiphonectes) hastatoides Fabricius, 1798

MoV sp. 5189

Records: 1 specimen, 21°59'S, 166 m

Distribution: Indo-West Pacific including Australia

Reference: Stephenson (1972) [photo below]

Portunus (Xiphonectes) longispinosus (Dana, 1852)

MoV sp. 5191

Records: 36 specimens, 22°50'S–24°01'S, 100 m

Distribution: Indo-West Pacific including N Australia; first record for WA – a species complex according to Davie

Reference: Davie (2002) [photo below]



Portunus aff. *orbitosinus* Rathbun, 1911

MoV sp. 5288

Records: 1 specimen, 21°58'S, 107 m

Distribution: new species? (det. V. Spiridonov) [photo next page]



Portunus nipponensis (Sakai, 1938)

MoV sp. 5126

Records: 1 specimen, 22°50'S, 100 m ✓

Distribution: Japan; first record for WA

Reference: Stephenson (1972) [photo below]



Portunus (Xiphonectes) pulchricristatus (Gordon, 1931)

MoV sp. 5184

Records: many specimens, 20°59'S–21°59'S, 100–166 m ✓

Distribution: Indo-West Pacific including NW Australia; first record for WA

Reference: Davie (2002) [photo upper right]



Thalamita macropus Montgomery, 1931

MoV sp. 5187

Records: 20 specimens, 21°59'S–33°58'S, 85–210 m ✓

Distribution: N Australia

Reference: Poore (2004) [photo below]



Thalamita sexlobata Miers, 1886

MoV sp. 5186

Records: 1 specimens, 27°48'S, 123–112 m

Distribution: Indo-West Pacific including Qld; first record for WA

Reference: Stephenson (1972) [photo below]



Thalamita spinifera Borradaile, 1902

MoV sp. 5291

Records: 3 specimens, 20°59'S–27°48'S, 100–123 m

Distribution: Indo-West Pacific including Cartier Reef; first record for S WA (det. V. Spiridonov)

Reference: Short and Davie (1993) [photo below]



Portunid sp. MoV 5289

Records: 5 specimens, 35°21'S, 91 m

Distribution: probable new genus and new species (det. V. Spiridonov) [photo below]



Superfamily Retroplumidae

Retroplumidae

A single species, doubtfully identified using de Saint Laurent (1989) is the first record of the family from Australia.

Retropluma* cf. *quadrata de Saint Laurent, 1989

MoV sp. 5093

Records: 1 specimen, 21°58'S, 373–382 m ✓

Distribution: if correctly identified, first Australian record for W Pacific species

Reference: de Saint Laurent (1989)

Superfamily Trapezioidea

Trapeziidae

The only species is recorded for the first time from Australia (Castro et al., 2004).

Quadrella reticulata Alcock, 1898

MoV sp. 5059

Records: 6 specimens, 21°57'S–27°48'S, 96–104 m ✓

Distribution: Indo-West Pacific; first record for Australia

Reference: Castro et al. (2004) [photo below]



Superfamily Xanthoidea

Panopeidae

A single species was identified using Ng's (1998) key to families and Davie (2002).

Homoioplax haswelli (Miers, 1884)

MoV sp. 5485

Records: 1 specimen, 21°58'S, 177–170 m

Distribution: Indo-West Pacific including N Australia; first record for S WA

Reference: Davie (2002)

Xanthidae

Eighteen species were found but proved difficult to identify using the standard text (Serène, 1984). With the help of Peter Davie, Queensland Museum, 13 taxa were identified to species or probable species using his unpublished notes. Of those that were identified to species, five are new Australian records.

Actaea calculosa (Milne Edwards, 1834)

MoV sp. 5116

Records: 1 specimen, 33°58'S, 96 m

Distribution: Australia

Reference: Poore (2004) [photo below]



Actaea peronii Milne Edwards, 1834

MoV sp. 1656

Records: 6 specimens, 33°02'S–35°20'S, 95–100 m

Distribution: Australia

Reference: Poore (2004) [photo below]



Atergatopsis cf. *alcocki* (Laurie, 1906)

MoV sp. 5117

Records: 2 specimens, 21°59'S, 166 m

Distribution: Indo-West Pacific including Qld; first record for Australia (det. P. Davie)

Reference: Sakai (1976) [photo below]



Calvactaea tumida Ward, 1933

MoV sp. 5083

Records: 1 specimen, 22°04'S, 106–101 m

Distribution: Indo-West Pacific including Australia

Reference: Poore (2004) [photo below]




Chlorodiella laevissima (Dana, 1852)

MoV sp. 5110

Records: 2 specimens, 27°03'S, 97 m

Distribution: Indo-West Pacific including Australia

Reference: Serène (1984: pl. 36D) [photo below]



Monodaeus tuberculidens (Rathbun, 1911)

MoV sp. 5075

Records: 3 specimens, 21°59'S–21°56'S, 132–166 m

Distribution: E Indian Ocean; first record for Australia (det. P. Davie) [photo below]



Demania wardi Garth & Ng, 1985

MoV sp. 5071

Records: 1 specimen, 21°59'S, 166 m

Distribution: West Pacific including Qld; first record for WA (det. P. Davie)

Reference: Davie (1989) [photo below]



Nanocassiope sp. MoV 5087

Records: 1 specimen, 31°43'S, 102 m

Distribution: new species (det. P. Davie) [photo below]



Medaeus sp. MoV 5081

Records: 1 specimen, 21°59'S, 166 m

Distribution: new species (det. P. Davie) [photo upper right]

Nanocassiope sp. MoV 5299

Records: many specimens, 20°59'S–31°43'S, 85–120 m ✓

Distribution: new species close to *N. alcocki* (Rathbun, 1902). Several colour morphs were separated in some stations but colour and morphology of the anterolateral carapace could not be correlated by Poore.

Reference: Serène (1984: 209 for *alcocki*)) [photo below]



Novactaea cf. *michaelseni* (Odhner, 1925)

MoV sp. 5074

Records: 9 specimens, 24°37'S–35°11'S, 97–1157 m

Distribution: WA (det. P. Davie) [photo below]



Palapedia pelsartensis (Serène, 1972)

MoV sp. 5219

Records: 1 specimens, 24°01'S, 100 m

Distribution: WA

Reference: Ng (1993) [photo upper right]



Palapedia valentini Ng, 1993

MoV sp. 5118

Records: 1 specimen, 27°48'S, 123–112 m

Distribution: Singapore; new record for Australia

Reference: Ng (1993) [photo below]



Paractaea rufopunctata (Milne Edwards, 1834)

MoV sp. 5073

Records: 1 specimen, 21°59'S, 166 m

Distribution: Indo-West Pacific, Atlantic including Australia (det. P. Davie) [photo next page]



Paractaea sp. MoV 5109

Records: 3 specimens, 29°48'S, 114 m

Distribution: new species (det. P. Davie) [photo below]



Paraxanthias aff. *pachydactylus* (Milne Edwards, 1867)

MoV sp. 5076

Records: 4 specimen, 27°55'S, 253 m ✓

Distribution: possible new species close to Indo-West Pacific-Australian species (det. P. Davie) [photo below]



Paraxanthodes cf. *cumatodes* (McGilchrist, 1905)

MoV sp. 5072

Records: 1 specimen, 28°59'S, 180–183 m

Distribution: new record for Australia if correctly identified (det. P. Davie) [photo below]



Platypodia cf. *semigranosa* (Heller, 1861)

MoV sp. 5082

Records: 1 specimen, 27°48'S, 98 m

Distribution: Indo-West Pacific including Qld; first record for WA (det. P. Davie)

Caridea – shrimps

Seventeen families were represented by 88 nominal species. Twenty (23%) are new species, 13 new records for Australia, 14 new records for WA and 17 new records for S WA. Caridean shrimps can be identified to family and genus using the keys of Holthuis (1993). More recent works apply for some families. Families are arranged alphabetically.

Alpheidae

Eighteen species were separated using the three papers on the Australian fauna by Banner and Banner (1973; 1975; 1982). *Alpheus* and *Synalpheus* are the dominant genera. Eleven species were identified to known species, all widely distributed in the Indo-West Pacific region and already known from WA. Seven species could not be identified because of insufficient material (listed here as new). The family is renowned for cryptic species and a difficult taxonomy.

Alpheopsis aff. *trispinosa* (Stimpson, 1861)

MoV sp. 5410

Records: 1 specimen, 33°58'S, 96 m

Distribution: new species slightly different from Indo-West Pacific-Australian species

Reference: Banner and Banner (1982) [photo below]



Alpheopsis sp. MoV 5407

Records: 1 specimen, 21°59'S, 166 m

Distribution: new species

Reference: Banner and Banner (1973)

Alpheopsis sp. MoV 5408

Records: 2 specimens, 22°50.55'S–31°37'S, 100–210 m

Distribution: new species

Reference: Banner and Banner (1973)

Alpheus alcyone De Man, 1902

MoV sp. 5419

Records: 1 specimen, 22°51'S, 100 m

Distribution: Indo-West Pacific including WA

Reference: Banner and Banner (1982: 110)

Alpheus hailstonei Coutière, 1905

MoV sp. 5420

Records: 34 specimens, 27°48'S–35°11'S, 95–210 m

Distribution: Indo-West Pacific, including WA

Reference: Banner and Banner (1982: 38) [photos upper right]



Alpheus paralcione Coutière, 1905

MoV sp. 5418

Records: 2 specimens, 20°59'S–25°54'S, 100 m

Distribution: Indo-West Pacific including WA

Reference: Banner and Banner (1982: 113)

Alpheus parasocialis Banner & Banner, 1982

MoV sp. 0722

Records: 2 specimens, 35°14'S, , 728–710 m

Distribution: Indo-West Pacific including WA

Reference: Banner and Banner (1982: 72)

Alpheus sp. MoV 5403

Records: 8 specimens, 22°50'S–25°54'S, 100 m ✓

Distribution: new species close to Australian *A. heronicus*

Reference: Banner and Banner (1982)



Alpheus sp. MoV 5405

Records: 2 specimens, 35°10'S, 97 m

Distribution: new species close to Australian *A. rapacida*

Reference: Banner and Banner (1982)

Alpheus sp. MoV 5406

Records: 13 specimens, 21°58'S–27°08'S, 373–414 m

Distribution: new species

Reference: Banner and Banner (1982)

***Alpheus* sp. MoV 5409**

Records: 8 specimens, 35°20'S–35°37'S, 99–179 m

Distribution: new species close to *A. distinguendus*

Reference: Banner and Banner (1982) [photo below]



***Synalpheus comatularum* (Haswell, 1882)**

MoV sp. 5413

Records: 5 specimens, 22°50'S–27°48'S, 96–100 m

Distribution: Indo-West Pacific including WA

Reference: Banner and Banner (1975) [photo below]



***Synalpheus lophodactylus* Coutière, 1908**

MoV sp. 5417

Records: 5 specimens, 28°58'S, 85 m

Distribution: Indo-West Pacific including WA

Reference: Banner and Banner (1975: 350)

***Synalpheus neomeris* (De Man, 1897)**

MoV sp. 5412

Records: 1 specimen, 27°48'S, 123–112 m

Distribution: Indo-West Pacific including WA

Reference: Banner and Banner (1975: 357)

***Synalpheus neptunus* (Dana, 1852)**

MoV sp. 5416

Records: 21 specimens, 21°59'S–33°58'S, 96–166 m ✓

Distribution: Indo-West Pacific including WA

Reference: Banner and Banner (1975: 317) [photo below]



***Synalpheus nilandensis* Coutière, 1905**

MoV sp. 5414

Records: 6 specimens, 22°37'S–25°54'S, 100–382 m

Distribution: Indo-West Pacific including WA

Reference: Banner and Banner (1975)

***Synalpheus streptodactylus* Coutière, 1905**

MoV sp. 5415

Records: 8 specimens, 21°58'S–27°48'S, 96–112 m

Distribution: Indo-West Pacific including WA (some taken from crinoid)

Reference: Banner and Banner (1975: 362)

***Synalpheus theano* De Man, 1910**

MoV sp. 5411

Records: 1 specimen, 27°03'S, 106 m

Distribution: Indo-West Pacific including WA

Reference: Banner and Banner (1975: 314)

Anchistioididae

The family was treated by Chace (1993). Our specimen was identified by A.J. Bruce.

Anchistioides willeyi (Borradaile, 1899)

MoV sp. 5424

Records: 1 specimen, 20°59'S, 100 m

Distribution: Indo-West Pacific including GBR, first record for WA

Reference: Chace (1993) ; det. A.J. Bruce

Bathypalaemonellidae

A single known species was recorded (Chace, 1997).

Bathypalaemonella pilosipes Bruce, 1986

MoV sp. 5449

Records: 5 specimens, 29°03'S, 1000–1037 m

Distribution: northern WA and Philippines, previously recorded to 400 m depth; new record for S WA

Reference: Chace (1997)

Bresiliidae

One species previously recorded from NSW was identified using Kensley (1983).

Discias brownae Kensley, 1983

MoV sp. 5428

Records: 1 specimen, 35°11'S, 157–147 m

Distribution: NSW; new record for WA

Reference: Kensley (1983)

Campylonotidae

A single well known southern species was recorded and identified from Poore (2004).

Campylonotus rathbunae Schmitt, 1926

MoV sp. 1806

Records: 5 specimens, 35°22'S–35°22'S, 676–728 m ✓

Distribution: southern Australian, New Zealand

Reference: Poore (2004) [photo below]



Crangonidae

The collection contains ten species of which six belong to known cosmopolitan or Indo-West Pacific species. None of the six are previously recorded from WA and only two from eastern Australia (Poore, 2004). The principal recent reference is by Chace (1984) and good illustrations appeared in De Man (1920). Uncertainty surrounds the specific and generic identification of some taxa.

Aegaeon lacazei (Gourret, 1887)

MoV sp. 1873

Records: 18 specimens, 24°33'S–31°00'S, 100–414 m ✓

Distribution: cosmopolitan; new record for WA

Reference: Chan (1996) [photo below]



Metacrangon sp. MoV 5423

Records: 16 specimens, 31°59'S–35°22'S, 408–728 m

Distribution: new species

Reference: Holthuis (1993) [photo below]



Parapontocaris aspera Chace, 1984

MoV sp. 5349

Records: 9 specimens, 21°58'S–22°04'S, 373–399 m

Distribution: Philippines; new Australian record

Reference: Chace (1984: 31)

Parapontocaris levigata Chace, 1984

MoV sp. 5350

Records: 22 specimens, 21°58'S–22°04'S, 324–399 m ✓

Distribution: Philippines; new Australian record

Reference: Chace (1984: 34) [photo upper right]



Parapontophilus junceus (Bate, 1888)

MoV sp. 5351

Records: 28 specimens, 22°00'S–35°22'S, 539–1077 m

Distribution: Indonesia-Philippines; new Australian record

Reference: Chace (1984: 53) [photo below]



Philocheras sp. MoV 5422

Records: 4 specimens, 21°58'S–25°55'S, 101–120 m ✓

Distribution: new species close to *P. magniculus*

Reference: Chace (1984); Komai and Chan (2007) [photo below]



Philocheras sp. MoV 5439

Records: 1 specimen, 23°59'S, 411 m

Distribution: new species

Reference: Chace (1984)

Pontocaris pennata Bate, 1888

MoV sp. 5353

Records: 5 specimens, 20°59'S–22°04'S, 102 m

Distribution: Indo-West Pacific, Indonesia; new Australian record

Reference: Chace (1984: 42); De Man (1920: pl. 24, fig. 70) [photo below]



Pontocaris propensalata Bate, 1888

MoV sp. 5354

Records: 2 specimens, 21°59'S, 166 m

Distribution: Philippines, Indonesia, NSW; new record for WA

Reference: Chace (1984: 43); De Man (1920: pl. 24, fig. 71)

Sabinea sp. MoV 5421

Records: 1 specimen, 33°00'S, 421–414 m

Distribution: like *S. indica* but carapace strongly depressed posterior to middorsal crest

Reference: De Man (1920: pl. 25, fig. 75) [photo below]



Eugatonotonotidae

The only species is previously recorded from NW Australia (Chace, 1997).

Eugonatonotus chacei Chan & Yu, 1991

MoV sp. 5429

Records: 1 specimen, 22°04'S, 399–387 m

Distribution: Eastern Pacific, NW Australia; new record for S WA

Reference: Chace (1997: 23) [photo below]



Glyphocrangonidae

Two described species known from Australia, one new Australian record, one newly recorded for S WA, and a fourth identified with some uncertainty to a Japanese species comprise the collection. Komai's recent paper (2004) is comprehensive.

Glyphocrangon lineata Komai, 2004

MoV sp. 5356

Records: 2 specimens, 21°58'S–22°00'S, 658–754 m ✓

Distribution: Indonesia, NW Australia; new record for S WA

Reference: Komai (2004) [photos below]



Glyphocrangon sibogae De Man, 1918

MoV sp. 5358

Records: 1 specimens, 21°55'S, 1260–1295 m ✓

Distribution: Indonesia, new Australian record

Reference: Komai (2004) [photos below]



Glyphocrangon cf. *perplexa* Komai, 2004

MoV sp. 5357

Records: 4 specimens, 21°58'S, 726–732 m, stn 159(4)

Distribution: probable new species similar to this Japanese species

Reference: Komai (2004)

Glyphocrangon confusa Komai, 2004

MoV sp. 5355

Records: 1 specimen, 29°00'S, 704–700 m

Distribution: Indonesia, NW Australia; new record for S WA

Reference: Komai (2004: 597) [photo below]



Hippolytidae

Five species were identified with reference to Poore (2004) and Chace (1997). Two were recorded for the first time from WA. Two are new species, one previously recorded from the Tasmanian seamounts.

Eualus sp. MoV 2681

Records: 5 specimens, 35°25'S–35°26'S, 900–980 m

Distribution: Tas. Seamounts; new species

Reference: Poore et al. (1998) [photo below]



Lebbeus sp. MoV 5425

Records: 1 specimen, 35°12'S, 431–408 m

Distribution: new species

Reference: Chace (1997) [photos below]




Lysmata amboinensis (De Man, 1888)

MoV sp. 5359

Records: 1 specimens, 22°50'S, 100 m

Distribution: Indo-West Pacific species; new record for S WA

Reference: Chace (1997) [photos below]



Merhippolyte chacei Kensley, Tranter & Griffin, 1987

MoV sp. 2615

Records: 5 specimens, 35°14'S–35°22'S, 676–728 m

Distribution: NSW, Tas., new WA record

Reference: Kensley et al. (1987) [photo below]



Tozeuma tomentosum (Baker, 1904)

MoV sp. 5361

Records: 16 specimens, 20°59'S–27°03'S, 100 m

Distribution: SA (doubtful record from Japan); new WA record

Reference: Chace (1997: 95) [photo below]



Nematocarcinidae

Three of four species of *Nematocarcinus*, previously recorded from WA, could be identified (Hanamura and Evans, 1996; Burukovskii, 2000). The fourth is an undescribed species recorded by Poore (2004).

Nematocarcinus hanamuri Burukovskii, 2000

MoV sp. 5452

Records: 3 specimens, 21°55'S, 1260–1295 m

Distribution: SW Australia

Reference: Burukovskii (2000)

Nematocarcinus productus Bate, 1888

MoV sp. 5450

Records: 2 specimens, 35°31'S–35°31'S, 1073–1110

Distribution: Indo-West Pacific, WA; new record for S WA

Reference: Hanamura and Evans (1996)

Nematocarcinus tenuirostris Bate, 1888

MoV sp. 5451

Records: 1 specimen, 21°55'S, 1260–1295 m

Distribution: Indo-West Pacific, WA

Reference: Hanamura and Evans (1996)

Nematocarcinus sp. MoV 5456

MoV sp. 5456

Records: 2 specimens, 35°16'S–35°31'S, 978–1110 m

Distribution: NSW, Tas.; new species that keys to *N. altus*

Reference: Poore (2004: fig. 17d) [photo below]



Oplophoridae

The seven species include one new record for Australia (Chace, 1986).

AcanthePHYra armata Milne Edwards, 1881

MoV sp. 5362

Records: 9 specimens, 21°58'S–22°00'S, 658–1010 m

Distribution: cosmopolitan, WA; new record for S WA

Reference: Wadley and Evans (1992: 13) [photo below]



AcanthePHYra faxoni Calman, 1939

MoV sp. 5430

Records: 3 specimens, 21°56'S–22°00'S, 1051–1077 m

Distribution: Indo-West Pacific; new Australian record

Reference: Chace (1986: key)

AcanthePHYra quadrispinosa Kemp, 1939

MoV sp. 1840

Records: 7 specimens, 21°55'S–35°04'S, 378–1295 m

Distribution: cosmopolitan, including WA

Reference: Wadley and Evans (1992) [photo below]



Janicella spinicauda (Milne Edwards, 1883)

MoV sp. 5431

Records: 2 specimens, 22°00'S, 983–1010 m

Distribution: cosmopolitan, including WA; new record for S WA

Reference: Hanamura (1987)

Oplophorus gracilirostris Milne Edwards, 1881

MoV sp. 5363

Records: 3 specimens, 21°58'S–22°50'S, 356–430 m ✓

Distribution: cosmopolitan, including WA

Reference: Hanamura (1987) [photo next page]



Oplophorus novaezeelandiae (De Man, 1931)

MoV sp. 1845

Records: 1 specimen, 35°16'S, 978–980 m

Distribution: cosmopolitan, including WA

Reference: Kensley et al.(1987) [photo below]



Systellaspis debilis (Milne Edwards, 1881)

MoV sp. 1841

Records: 3 specimens, 22°00'S–22°00S, 983–1085 m

Distribution: cosmopolitan, including WA

Reference: Kensley et al. (1987), Poore (2004)

Palaemonidae

Although common in shallow waters this family was represented by only three specimens. The identifications below are by A.J. Bruce.

Periclimenes aleator Bruce, 1991

MoV sp. 5448

Records: 1 specimen, 21°00'S, 399–408 m

Distribution: Loyalty Is., new record for Australia

Reference: Bruce (1991); det. A.J. Bruce

Palaemonid sp. MoV 5437

Records: 2 specimens, 29°48'S, 114 m

Distribution: new genus and species

Reference: det. A.J. Bruce

Pandalidae

Of 21 species of mostly benthopelagic shrimps, 15 are recorded outside their known range. Three are new Australian records of Indo-West Pacific species and four probable new species. Four studies have covered the family in this region (Chace, 1985; Hanamura and Takeda, 1987; Crosnier, 1988; Hanamura and Evans, 1996).

Chlorotocella spinicaudus (Milne Edwards, 1837)

MoV sp. 0995

Records: 1 specimen, 23°59'S, 411 m

Distribution: common southern Australian species

Reference: Poore (2004: 131)

Chlorotocus sp. MoV 5443

Records: 1 specimen, 21°58'S, 356–324 m ✓

Distribution: new species

References: Hanamura and Takeda (1987); Hanamura and Evans (1996)

Heterocarpoides levicarina Bate, 1888

MoV sp. 5364

Records: 14 specimens, 21°58'S–22°04'S, 101–206 m

Distribution: Indo-West Pacific, including Indonesia; new Australian record

Reference: Chace (1985: 17)

Heterocarpus dorsalis Bate, 1888

MoV sp. 5365

Records: 25 specimens, 21°55'S–31°57'S, 726–1260 m ✓

Distribution: cosmopolitan, including Australia

Reference: Wadley and Evans (1992) [photos below]



Heterocarpus hayashii Crosnier, 1988

MoV sp. 5541

Records: 16 specimens, 21°58'S–27°08'S, 373–431 m

Distribution: West Pacific, including GBR, Australia; new record for WA

Reference: Crosnier (1988) [photo upper right]



Heterocarpus tricarinatus Alcock & Anderson, 1894

MoV sp. 5366

Records: 5 specimens, 21°55'S, 1260–1295 m

Distribution: Indo-West Pacific, including N WA; new record for S WA

Reference: Hanamura and Evans (1996: 9) [photo below]



Heterocarpus woodmasoni Alcock, 1901

MoV sp. 5367

Records: 18 specimens, 21°58'S–22°50'S, 373–430 m

Distribution: Indo-West Pacific, including N Australia; new record for S WA

Reference: Hanamura and Evans (1996: 10) [photo below]



Heterocarpus MoV sp. 5540

Records: 25 specimens, 21°00'S–22°04'S, 399–411 m

Distribution: new species

Reference: Crosnier (1988) [photo below]



Plesionika bifurca Alcock & Anderson, 1894

MoV sp. 5444

Records: 2 specimens, 22°00'S, 983–1010 m

Distribution: Indo-West Pacific, including N Australia; first record for S WA

Reference: Hanamura and Takeda (1987); Chace (1985)

Plesionika binoculus (Bate, 1888)

MoV sp. 5447

Records: 6 specimens, 21°59'S, 166 m

Distribution: Arafura Sea; first record for S WA

Reference: Chace (1985: key)

Plesionika* cf. *kensleyi Chace, 1985

MoV sp. 5369

Records: 33 specimens, 22°04'S–35°14'S, 212–1050 m

Distribution: rostrum more compact than *P. kensleyi*; new Australian record or new species

Reference: Chace (1985: 77) [photo below]



Plesionika* cf. *philippinensis Chace, 1985

MoV sp. 5370

Records: 1 specimen, 21°57'S, 104–114 m

Distribution: new Australian record or new species

Reference: Chace (1985: 99)

Plesionika edwardsii (Brandt, 1851)

MoV sp. 5368

Records: 6 specimens, 24°33'S–31°55'S, 364–484 m ✓

Distribution: cosmopolitan; new record for WA

Reference: Chace (1985: 62)

Plesionika orientalis Chace, 1985

MoV sp. 5445

Records: many specimens, 21°58'S–35°12'S, 324–695 m ✓

Distribution: Indo-West Pacific usually as *P. martia orientalis*; first record for S WA

Reference: Chace (1985: 62) [photo below]



Plesionika reflexa Chace, 1985

MoV sp. 5371

Records: 40 specimens, 21°00'S, 399–408 m

Distribution: Indo-West Pacific, including N Australia; first record for S WA

Reference: Hanamura and Takeda (1987)

Plesionika semilaevis Bate, 1888

MoV sp. 5372

Records: 12 specimens, 22°04'S–35°22'S, 387–680 m ✓

Distribution: Indo-West Pacific, including N Australia; first record for S WA

Reference: Chace (1985: 113)

Plesionika serratifrons Borradaile, 1899

MoV sp. 5373

Records: many specimens, 21°58'S–27°55'S, 100–253 m ✓

Distribution: Indo-West Pacific; new record for Australia

Reference: Chace (1985: 123) [photos below]



Plesionika spinidorsalis (Rathbun, 1906)

MoV sp. 5374

Records: 23 specimens, 21°58'S–22°04'S, 356–754 m ✓

Distribution: Indo-West Pacific, including N WA; new record for S WA

Reference: Hanamura and Takeda (1987); Chace (1985)

Plesionika spinipes Bate, 1888

MoV sp. 5446

Records: 3 specimens, 22°04'S, 106–101 m

Distribution: Indo-West Pacific, including E Australia; new record for WA

Reference: Chace (1985: key)

***Plesionika* sp. MoV 5457**

Records: 1 specimen, 35°16'S, 980–976 m

Distribution: probable new species

Reference: Chace (1985)

Procletes levicarina (Bate, 1888)

MoV sp. 5483

Records: 2 specimens, 21°59'S, 166 m

Distribution: Indo-West Pacific, including N Australia; new record for S WA

Reference: Holthuis (1993)

Pasiphaeidae

The six species included three already known from WA, two new WA records and a probable new species. Hanamura & Evans (1994) is a key reference.

Alainopasiphaea australis (Hanamura, 1989)

MoV sp. 1895

Records: 2 specimens, 35°22'S, 676–680 m

Distribution: southern Australia

References: Hayashi (2004), Poore (2004); Hanamura (1989)

Eupasiphae sp. MoV 5427

Records: 2 specimens, 21°56'S–21°58'S, 726–1050 m

Distribution: new species

Reference: Hanamura and Evans (1994)

Leptochela sydniensis Dakin & Colefax, 1940

MoV sp. 0723

Records: 5 specimens, 22°04'S–35°18'S, 95–210 m

Distribution: Indo-West Pacific, including N, E and S Australian coasts; first record for WA

Reference: Hanamura and Evans (1994) [photo below]



Pasiphaea kapala Kensley, Tranter & Griffin, 1987

MoV sp. 5432

Records: 2 specimens, 35°22'S, 685–695 m

Distribution: southern Australia

Reference: Poore (2004)

Pasiphaea longitaenia Kensley, Tranter & Griffin, 1987

MoV sp. 5377

Records: 1 specimen, 22°00'S, 983–1010 m

Distribution: NSW; new record for WA

Reference: Kensley et al.(1987)

Pasiphaea tarda Krøyer, 1845

MoV sp. 5433

Records: 1 specimen, 35°31'S, 1074–1080 m

Distribution: cosmopolitan, including S WA

Reference: Hanamura and Evans (1994)

Processidae

Chace (1997) is the key reference. The three species included one known previously from WA, and two newly recorded for Australia. None has been adequately figured recently.

Hayashidonus japonicus (De Haan, 1844)

MoV sp. 5434

Records: 2 specimens, 21°59'S–22°04'S, 101–166 m

Distribution: Indo-West Pacific including Indonesia; new record for Australia

Reference: Chace (1997: 33)

Processa gracilis Baker, 1907

MoV sp. 5376

Records: 1 specimen, 33°58'S, 96 m

Distribution: SA; first record for WA

Reference: Poore (2004: 128)

Processa longirostris Hayashi, 1975

MoV sp. 5426

Records: 8 specimens, 21°58'S–24°01'S, 100–107 m ✓

Distribution: S Vietnam; new Australian record

References: Hayashi (1975: key); Noël (1986: key)

Rhynchocinetidae

Two described species, one known from southern Australia and the other from northern Australia, were found and identified using Okuno (1994) and Chace (1997).

Rhynchocinetes brucei Okuno, 1994

MoV sp. 5378

Records: 17 specimens, 21°59'S–35°13'S, 100–494 m ✓

Distribution: West Pacific, NE Australia; new record for WA

Reference: Okuno (1994) [photo below]



Rhynchocinetes enigma Okuno, 1997

MoV sp. 3978

Records: 19 specimens, 31°37'S–35°21'S, 97–210 m

Distribution: S Australia; new record for WA

Reference: Poore (2004: 76) [photo below]



Thalassocarididae

One Indo-West Pacific species was recorded for the first time from Australia (Chace, 1985).

Thalassocaris crinita (Dana, 1852)

MoV sp. 5379

Records: 6 specimens, 22°50'S–27°03'S, 97–100 m

Distribution: Indo-West Pacific; first record for Australia

Reference: Chace (1985: 7)

Polychelida – deep sea lobsters

The Australian fauna is well studied and two papers enabled the collections (59 individuals) to be identified (Galil, 2000; Ahyong and Brown, 2002).

Polychelidae

All five species in two genera are already described. *Polycheles coccifer* Galil, 2000 was previously recorded from Indonesia so this record from northerly stations is not unexpected.

Pentacheles laevis Bate, 1878

MoV sp. 3980

Records: 5 specimens, 31°57'S–35°31'S, 928–1170 m

Distribution: cosmopolitan species, including S Australia

References: Ahyong and Brown (2002) [photo below]



Polycheles auriculatus (Bate, 1878)

MoV sp. 4975

Records: 36 specimens, 21°56'S–35°14'S, 658–1037 m ✓

Distribution: Indo-West Pacific species, including WA

References: Ahyong and Brown (2002) [photo below]



Polycheles coccifer Galil, 2000

MoV sp. 4973

Records: 3 specimens, 21°58'S–21°58'S, 324–382 m ✓

Distribution: Indo-West Pacific; first record for Australia

References: Galil (2000) [photos below]



Polycheles suhmi (Bate, 1878)

MoV sp. 3979

Records: 2 specimens, 35°14'S–35°22'S, 676–728 m

Distribution: Southern Ocean, including NSW–Tas.; first record for WA

References: Galil (2000) [photo below]



Polycheles typhlops Heller, 1862

MoV sp. 5069

Records: 3 specimens, 21°58'S–22°04'S, 373–399 m

Distribution: cosmopolitan species

References: figures from Galil (2000)

Stenopodidea – coral shrimps

Stenopodidean shrimps can be identified to family and genus using the keys of Holthuis (1993). One species was represented by one individual of a species previously recorded from the region but not so far south. The other was just a cheliped but could be identified as probably a species not recorded from Australia.

Stenopodidae

Two species were found, one known from coral in Indonesia and northern WA and the other, represented in this collection by a single cheliped, from throughout the Indo-West Pacific.

Engystenopus cf. *palmipes* Alcock & Anderson, 1894

MoV sp. 5545

Records: 1 detached cheliped (pereopod 3), 22°04'S, 400 m ✓

Distribution: Bay of Bengal, Philippines; new record for Australia (det. J. Goy from photo of cheliped)

References: De Saint Laurent and Cleva (1981) [photo below]



Odontozona sculpticaudata Holthuis, 1946

MoV sp. 5442

Records: 1 specimen, 22°50'S, 100 m

Distribution: Indo-West Pacific species, including N Australia; new record for S WA

References: Holthuis (1946)

Thalassinidea – ghost and sponge shrimps

Six families (of 11 known) are represented by 23 species. The collection is not large, 51 individuals of which 13 belonged in one species. Surprisingly, only four species could be identified, one of these with a Korean species newly recorded from Australia. The fraction of new species is 82%. Several seemed not to fit well with presently diagnosed genera. Published keys to families and genera (Poore, 1994) are now superseded by an interactive DELTA-based key in preparation by Poore. Poore and Griffin (1979) covered all the Australian species then known but as citations below indicate, the number has grown since.

Axiidae

Of the eight species, at least one is most probably a new genus. None belong in the taxa described by Sakai (1986; 1994) or Kensley (1989). One is tentatively identified as a species described from Korea.

Acanthaxius sp. MoV 4956

Records: 1 specimen, 21°59'S, 166 m ✓

Distribution: new species

Reference: Ngoc-Ho (2006) [photo below]



Axiopsis tsushimaensis Sakai, 1992

MoV sp. 5440

Records: 2 specimens, 29°48'S–35°11'S, 113–157 m

Distribution: Korea, Japan; new record for Australia

Reference: Sakai (1992)

Axiopsis sp. MoV 5435

Records: 2 specimens, 27°48'S, 96–98 m

Distribution: new species

Reference: Poore (1994)

***Bouvieraxius* sp. MoV 4959**

Records: 1 specimen, 27°08'S, 414–405 m

Distribution: new species

Reference: Poore (1994) [photo below]



***Calocarides* sp. MoV 4955**

Records: 3 specimens, 25°54'S–27°03'S, 97–100 m

Distribution: new species

Reference: Poore (1994) [photos below]



***Calocarides* sp. MoV 4957**

Records: 1 specimen, 25°55'S, 404–407 m

Distribution: new species

Reference: Poore (1994) [photo upper right]



***Dorphinaxius* sp. MoV 4958**

Records: 1 specimen, 20°59'S, 100 m ✓

Distribution: new species

Reference: Poore (1994)

***Marianaxius* sp. MoV 5436**

Records: 1 specimens, 29°48'S, 114 m

Distribution: new species

Reference: Kensley (2003)

***Axiid* sp. MoV 5527**

Records: 1 specimens, Station not recorded

Distribution: new species, genus indetermined

***Axiid* sp. MoV 4954**

Records: 1 specimen, 33°00'S, 423–397 m

Distribution: new species, possibly new genus

Reference: Poore (1994) [photo below]



Callianassidae

There are only one or two individuals of each of the six species, sometimes incomplete as is typical of members of this family. One species has already been described in a manuscript in press (Poore, in press) but the others are not in papers dealing with the fauna of this region (Sakai, 1988; Ngoc-Ho, 1994; Poore, in press). Generic concepts in Callianassidae are unclear – most could not be placed in one of the 20 genera diagnosed in a DELTA key currently under construction. The most recent reviews of the family (Sakai, 1999, 2005) proposed an idiosyncratic taxonomy that does not recognise many traditionally recognised genera. For the time being, most species in this collection are tentatively placed in the catch-all ‘*Callianassa*’.

Callianassa sp. MoV 4964

Records: 1 specimens, 22°50’S, 100 m

Distribution: WA

Reference: Poore (in press)

Callianassa sp. MoV 4961

Records: 2 specimens, 22°04’S, 206–201 m

Distribution: new species

Callianassa sp. MoV 4962

Records: 1 specimen, 22°04’S, 206–201 m

Distribution: new species

Callianassa sp. MoV 4963

Records: 1 specimen, 29°48’S, 114 m

Distribution: new species

Callianassa sp. MoV 4966

Records: 1 specimen, 21°58’S, 107 m

Distribution: new species

Corallianassa sp. MoV 4965

Records: 2 specimens, 31°43’S–35°11’S, 102–169 m

Distribution: new species [photos below]



Calocarididae

The single species belongs to a genus not previously recorded from Australia.

Ambiaxius sp. MoV 4967

Records: 2 specimens, 33°00’S, 423–397 m

Distribution: new species

Reference: Sakai and Ohta (2005)

Gourretiidae

One new species was found whose generic identification is problematic. The nomenclature, composition and definition of this family is subject to considerable debate. The views of Sakai (2005) who provided the most recent revision are not necessarily followed here (see too Callianassidae).

Lipkecallianassa sp. MoV 4960

Records: 8 specimens, 21°59'S–22°04'S, 100–206 m

Distribution: generic placement of the species is problematic

Reference: Sakai (2002)

Micheleidae

The single specimen in each of two genera does not belong to any of the Western Australian (or other) species described by Poore (1997; in press).

Michelea sp. MoV 4969

Records: 1 specimen, 27°48'S, 123–112 m

Distribution: new species

Reference: Poore (1997)

Tethisea sp. MoV 5472

Records: 1 specimen, 35°22'S, 419–460 m

Distribution: new species, possibly new genus

Reference: Poore (1997)

Upogebiidae

All three species were identified by N. Ngoc-Ho and have been previously recorded from Australia.

Upogebia ancylodactyla De Man, 1905

MoV sp. 5078

Records: 2 specimens, 31°43'S, 102 m

Distribution: Indonesia–Philippines, N Australia; new record for S WA

Reference: Sakai (1993) [photo below]



Upogebia holthuisi Sakai, 1982

MoV sp. 4970

Records: 2 specimens, 25°54'S, 100 m

Distribution: New Caledonia, Pacific, first record for Australia

Reference: Sakai (1982)

Upogebia bowerbanki (Miers, 1884)

MoV sp. 4971

Records: 13 specimens, 21°57'S, 104–114 m ✓

Distribution: S Australia

Reference: Poore (2004) [photos below]



Acknowledgements

We acknowledge the considerable investment and enthusiasm of our colleagues at CSIRO Marine and Atmospheric Research (CMAR). Alan Williams and Rudy Kloser, co-principal investigators of the “Voyages of Discovery” research program were largely responsible for the sampling design and for securing the funding that made it happen. We thank these two and their CMAR colleagues Mark Lewis (Gear Officer) and Karen Gowlett-Holmes (Curator, Marine Invertebrates) for help with gear and dealing with samples while on board FRV *Southern Surveyor*. Special thanks to Karen Gowlett-Holmes for the many photographs of live animals taken on board and published here.

We appreciate the assistance of the shipboard invertebrate zoology crew – Tim O’Hara, David Staples, Anna Syme, Robin Wilson (Museum Victoria), Jane Fromont, Mark Salotti, Corey Whisson, Shirley Slack-Smith (Western Australian Museum), Penny Berents and Steve Keable (Australian Museum) – for their tireless dedication to sorting the catches. We thank the master and crew of FRV *Southern Surveyor* for their cooperation.

In the lab in Melbourne we must thank the student volunteers who assisted with the unpacking, preliminary sorting and labelling – Amanda Schaarschmidt, Sarah Thompson and Kirsteen Roberts. We thank too Kate Naughton whose first sort of hermit crabs and shrimps made our job easier.

Most identifications were done at Museum Victoria but we sought the help of Peter Davie, Queensland Museum, for the enigmatic pilumnid and xanthoid crabs, Nguyen Ngoc-Ho, Muséum nationale d’Histoire naturelle, Paris, for upogebiid shrimps, and Joseph Goy, Harding University, Arkansas, for one stenopidean shrimp claw – for which we are extremely grateful. Vasily Spiridonov provided identifications of some portunid crabs, Bertrand Richer de Forges re-examined some majoid crabs, and Sandy Bruce our few palaemonid and anchistioid shrimps.

For financial support we acknowledge the Commonwealth Department of the Environment and the CSIRO Wealth from Oceans Flagship that provided funds for the field and laboratory components of the “Voyages of Discovery” program. Thanks too to OBIS Australia (Oceanographic Biological Information System) for support to enable the data to made publicly available on the OBIS website.

References

- Ahyong, S. 2008. Deepwater crabs from seamounts and chemosynthetic habitats off eastern New Zealand (Crustacea: Decapoda: Brachyura). *Zootaxa* 1708: 1-72.
- Ahyong, S., and Brown, D.E. 2002. New species and new records of Polychelidae from Australia (Crustacea: Decapoda). *Raffles Bulletin Of Zoology* 50: 53-79.
- Ahyong, S., and Brown, D.E. 2003. New species of *Cymonomus* from southeastern Australia (Brachyura, Cymonomidae) with a key to the Indo-West Pacific species. *Crustaceana* 75: 1363-1374.
- Ahyong, S., and O’Meally, D. 2004. Phylogeny of the Decapoda Reptantia: resolution using three molecular loci and morphology. *Raffles Bulletin Of Zoology* 52: 673-693.
- Ahyong, S.T., and Poore, G.C.B. 2004a. The Chirostylidae of southern Australia (Crustacea: Decapoda: Anomura). *Zootaxa* 436: 1-88.
- Ahyong, S.T., and Poore, G.C.B. 2004b. Deep-water Galatheidae (Crustacea: Decapoda: Anomura) from southern and eastern Australia. *Zootaxa* 472: 1-76.
- Ahyong, S.T., and Poore, G.C.B. 2004c. Deep-water Galatheidae (Crustacea : Decapoda : Anomura) from southern and eastern Australia. *Zootaxa* 472: 3-76.
- Alcock, A. 1895. Materials for a carcinological fauna of India. No. 1. The Brachyura Oxyrhyncha. *Journal of the Asiatic Society of Bengal* 64: 157-291, pls 153-155.
- Alcock, A., and Anderson, A.R.S. 1895. *Crustacea, Part III. Illustrations of the Zoology of the Royal Indian Marine Surveying Steamer Investigator, under the Command of Commander A. Carpenter, R.N., D.S.O., of the Late Commander R.F. Hoskyn, R.N., and of Commander C.F. Oldham*. Trustees of the Indian Museum: Calcutta. pls 9-15.
- Arnold, P., and George, R.W. 1987. Recognition of *Leucosia whitei* Bell and *Leucosia cheverti* Haswell (Decapoda, Brachyura, Leucosiidae). *Crustaceana* 53: 209-214.
- Asakura, A. 2001. A revision of the hermit crabs of the genera *Catapagurus* A. Milne-Edwards and *Hemipagurus* Smith from the Indo-West Pacific (Crustacea : Decapoda : Anomura : Paguridae) *Invertebrate Taxonomy* 15: 823 - 891.
- Baba, K. 1988. Chirostylid and galatheid crustaceans (Decapoda: Anomura) of the "Albatross" Philippine Expedition, 1907-1910. *Researches on Crustacea, Special Number 2*: 1-203.
- Baba, K. 1994. Deep-sea galatheid crustaceans (Anomura: Galatheidae) collected by the 'Cidarid I' Expedition off central Queensland, Australia. *Memoirs of the Queensland Museum* 35: 1-21.
- Baba, K. 2005. Deep-sea chirostylid and galatheid crustaceans (Decapoda: Anomura) from the Indo-West Pacific, with a list of species. *Galathea Reports* 20: 1-317.
- Baba, K., and Poore, G.C.B. 2002. *Munidopsis* (Decapoda, Anomura) from south-eastern Australia. *Crustaceana* 75: 231-252.
- Baker, W.H. 1905. Notes on South Australian decapod Crustacea. Part II. *Transactions of the Royal Society of South Australia* 29: 116-130, pls 121-124.
- Balss, H. 1935. Brachyura of the Hamburg Museum expedition to south-western Australia, 1905. *Journal of the Royal Society of Western Australia* 21: 113-151.
- Banner, D.M., and Banner, A.H. 1973. The alpheid shrimp of Australia. Part I: the lower genera. *Records of the Australian Museum* 28: 291-382.
- Banner, D.M., and Banner, A.H. 1975. The alpheid shrimp of Australia. Part 2: the genus *Synalpheus*. *Records of the Australian Museum* 29: 267-389.
- Banner, D.M., and Banner, A.H. 1982. The alpheid shrimp of Australia. Part III: the remaining alpheids, principally the genus *Alpheus*, and the family Ogyrididae. *Records of the Australian Museum* 34: 1-358.
- Barnard, K.H. 1950. Descriptive catalogue of South African decapod Crustacea (crabs and shrimps). *Annals of the South African Museum* 38: 1-837.
- Boyko, C.B. 2002. A worldwide revision of the Recent and fossil sand crabs of the Albuneidae Stimpson and Blepharipodidae, new family (Crustacea: Decapoda: Anomura: Hippoidea). *Bulletin of the American Museum of Natural History* 272: 1-396.
- Bruce, A.J. 1991. Shallow-water palaemonoid shrimps from New Caledonia (Crustacea: Decapoda). Pp. 221-279 in: Richer de Forges, B. (ed.) *Le benthos des fonds meubles*

- des lagons de Nouvelle-Calédonie*. Vol. 1. ORSTOM Editions: Paris.
- Burukovskii, R.N. 2000. Taxonomy of shrimps of the genus *Nematocarcinus* (Crustacea, Decapod, Nematocarcinidae). 7. Description of new species, *N. hanamuri* and *N. evansi*, from southwestern Australian waters. *Zoologicheskii Zhurnal, Moscow* 79: 1290-1293.
- Campbell, B.M., and Stephenson, W. 1970. The sublittoral Brachyura (Crustacea: Decapoda) of Moreton Bay. *Memoirs of the Queensland Museum* 15: 235-301, pl.222.
- Castro, P. 2000. Crustacea Decapoda: a revision of the Indo-West Pacific species of palicid crabs (Brachyura Palicidae). In: Crosnier, A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 21. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 184: 437-610.
- Castro, P. 2007. A reappraisal of the family Goneplacidae MacLeay, 1838 (Crustacea, Decapoda, Brachyura) and revision of the subfamily Goneplacinae, with the description of 10 new genera and 18 new species. *Zootaxa* 29: 609-774.
- Castro, P., and Davie, P.J.F. 2003. New records of palicid crabs (Crustacea: Brachyura: Palicidae) from Australia. *Memoirs of the Queensland Museum* 49: 153-157.
- Castro, P., Williams, A.B., and Cooper, L.L. 2003. Revision of the family Latreilliidae Stimpson, 1858 (Crustacea, Decapoda, Brachyura). *Zoosystema* 25: 601-634.
- Castro, P., Ng, P.K.L., and Ahyong, S. 2004. Phylogeny and systematics of the Trapeziidae Miers, 1886 (Crustacea: Brachyura), with the description of a new family. *Zootaxa* 643: 1-70.
- Chace, F.A. 1984. The caridean shrimps (Crustacea; Decapoda) of the Albatross Philippine Expedition, 1907-1910, part 2: families Glyphocrangonidae and Crangonidae. *Smithsonian Contributions to Zoology* 397: 1-63.
- Chace, F.A. 1985. The caridean shrimps (Crustacea; Decapoda) of the Albatross Philippine Expedition, 1907-1910, part 3: families Thalassocarididae and Pandalidae. *Smithsonian Contributions to Zoology* 411: 1-143.
- Chace, F.A. 1986. The caridean shrimps (Crustacea; Decapoda) of the Albatross Philippine Expedition, 1907-1910, part 4: families Oplophoridae and Nematocarcinidae. *Smithsonian Contributions to Zoology* 432: 1-82.
- Chace, F.A. 1997. The caridean shrimps (Crustacea: Decapoda) of the Albatross Philippine Expedition, 1907-1910, Part 7: Families Atyidae, Eragonotidae, Rhynchocinetidae, Bathypalaemonellidae, Processidae, and Hippolytidae. *Smithsonian Contributions to Zoology* 587: i-v, 1-106.
- Chace, F.A., and Bruce, A.J. 1993. The caridean shrimps (Crustacea; Decapoda) of the Albatross Philippine Expedition, 1907-1910, part 6: superfamily Palaemonoidea. *Smithsonian Contributions to Zoology* 543: 1-152.
- Chan, T.-Y. 1996. Crustacea Decapoda Crangonidae: revision of the three closely related genera *Aegaeon* Agassiz, 1846, *Pontocaris* Bate, 1888 and *Parapontocaris* Alcock, 1901. In: Crosnier, A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 15. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 168: 269-336.
- Chen, H. 1989. Leucosiidae (Crustacea, Brachyura). In: Forest, J. (ed.), Résultats des Campagnes MUSORSTOM, vol. 5. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 144: 181-264.
- Chen, H.L. 1993. Crustacea Decapoda: Dorippidae of New Caledonia, Indonesia and the Philippines. In: Crosnier, A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 10. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 156: 315-345.
- Chia, D.G.B., and Ng, P.K.L. 2000. A revision of *Eumedon* H. Milne Edwards, 1834 and *Gonatonotus* White, 1847 (Crustacea: Decapoda: Brachyura: Eumedonidae), two genera of crabs symbiotic with sea urchins. *Journal of Natural History* 34: 15-56.
- Clark, P.F., and Ng, P.K.L. 2006. A new species of *Notonyx* A. Milne-Edwards, 1873 (Crustacea, Brachyura, Goneplacidae) from the intertidal zone of Phuket, Thailand. *Zoosystema* 28: 539-551.
- Crosnier, A. 1985. Crustacés Décapodes: Penaeidae. Les espèces indo-ouest-pacifique du genre *Parapeneaeus*. In: Crosnier, A. (ed.), Résultats des Campagnes MUSORSTOM I et II - Philippines (1976, 1980). Tome 2. *Mémoires du Muséum National d'Histoire Naturelle, Paris, ser. A (Zoologie)* 133: 303-355.
- Crosnier, A. 1988. Sur les *Heterocarpus* (Crustacea, Decapoda, Pandalidae) du sud-ouest de l'océan Indien. Remarques sur d'autres espèces ouest-pacifiques du genre et description de quatre taxa nouveaux. *Bulletin du Muséum National d'Histoire Naturelle, Paris* 10: 57-103.
- Crosnier, A. 1991. Crustacea Decapoda: Les *Metapeneaeopsis* indo-ouest-pacifiques sans appareil strident (Penaeidae). Deuxième partie. In: Crosnier, A. (ed.), Résultats des Campagnes MUSORSTOM. Vol. 9. *Mémoires du Muséum National d'Histoire Naturelle, Paris, ser. A (Zoologie)* 152: 155-297.
- Crosnier, A. 2003. *Sicyonia* (Crustacea, Decapoda, Penaeoidea, Sicyoniidae) de l'indo-ouest Pacifique. *Zoosystema* 25: 197-350.
- Dall, W. 1999. Australian species of Solenoceridae (Penaeoidea: Decapoda). *Memoirs of the Queensland Museum* 43: 553-587.
- Dall, W. 2001. Australian species of Aristeidae and Benthescymidae (Penaeoidea: Decapoda). *Memoirs of the Queensland Museum* 46: 409-441.
- Dampier, W. 1703. *A voyage to New Holland, &c. in the year, 1699. Wherein are described, The Canary Islands, the Isles of Mayo and St. Jago. The Bay of All-Saints, with the forts and town of Bahia in Brazil. Cape Salvador. The winds on the Brazilian coast. Abrolho Shoals. A table of all the variations observed in this voyage. Occurrences near the Cape of Good Hope. The course to New Holland. Shark's Bay. The isles and coast, &c. of New Holland. Their inhabitants, manners, customs, trade, etc. Their harbours, soil, beasts, birds, fish, etc. Trees, plants, fruits, &c. Illustrated with several maps and draughts: also divers birds, fishes and plants not found in this part of the world, curiously engraven on copper plates.* James and John Knapton London.
- <http://www.canadiana.org/ECO/ItemRecord?id=e6263c2f9774a7b4>
- Davie, P.J.F. 1989a. New records of *Dermania* (Crustacea: Decapoda: Xanthidae) from Australia. *Memoirs of the Queensland Museum* 27: 123-128.
- Davie, P.J.F. 1989b. Two new genera of the family Pilumnidae (Crustacea: Decapoda: Brachyura) from Queensland, Australia. *Journal of Natural History* 23: 1353-1365.
- Davie, P.J.F. 2002a. *Crustacea: Malacostraca: Phyllocarida, Hoplocarida, Eucarida (Part 1)*. Vol. 19.3A. CSIRO Publishing: Melbourne. xii, 551 pp.

- Davie, P.J.F. 2002b. *Crustacea: Malacostraca: Eucarida (Part 2): Decapoda - Anomura, Brachyura*. Vol. 19.3B. CSIRO Publishing: Melbourne. xiv, 641 pp.
- Davie, P.J.F., and Turner, P.A. 1994. A new record and new species of *Parthenope* from northern Australia (Crustacea: Brachyura: Parthenopidae). *Raffles Bulletin Of Zoology* 42: 975-981.
- Davie, P.J.F., and Humpherys, A. 1997. New species of rhizopine crabs (Crustacea: Brachyura) from northern Australia. *Memoirs of the Queensland Museum* 42: 97-103.
- Davie, P.J.F., and Crosnier, A. 2006. *Echinolatus* n. gen. (Crustacea, Decapoda, Portunidae) with description of two new species from the South-West Pacific. In: Richer de Forges, B., and Justine, J.-L. (eds), *Tropical Deep-Sea Benthos*, Vol. 24. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 193: 393-410.
- Dawson, E.W., and Yaldwyn, J.C. 2000. Description and ecological distribution of a new frog crab (Crustacea, Brachyura, Raninidae) from northern New Zealand waters, with keys to Recent raninid genera and *Notosceles* species. *Tuhinga* 11: 47-71.
- Dell, R.K. 1968. A new crab of the genus *Trichopeltarion* from Australia. *Australian Zoologist* 14: 275-276, pl. 216.
- Flipse, H.J. 1930. Decapoda F. (Decapoda Brachyura continued) Oxyrhyncha: Parthenopidae. *Siboga Expédition* 39c: 1-96.
- Forest, J. 1987. Les Pylochelidae ou "Pagures symétriques" (Crustacea Coenobitoidea). *Mémoires du Muséum National d'Histoire Naturelle, Paris* 137: 1-254, 259 pls.
- Forest, J. 1995. Crustacea Decapoda Anomura: révision du genre *Trizopagurus* Forest, 1952 (Diogenidae), avec l'établissement de deux genres nouveaux. In: Crosnier, A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 13. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 163: 9-149.
- Galil, B.S. 1993. Crustacea Decapoda: a revision of the genus *Mursia* Desmarest, 1823 (Calappidae). In: Crosnier, A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 10. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 156: 347-379.
- Galil, B.S. 1997. Crustacea Decapoda: a revision of the Indo-Pacific species of the genus *Calappa* Weber, 1795 (Calappidae). In: Crosnier, A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 18. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 176: 271-335.
- Galil, B.S. 2000. Crustacea Decapoda: review of the genera and species of the family Polychelidae Wood-Mason, 1874. In: Crosnier, A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 21. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 184: 285-387.
- Galil, B.S. 2001a. A revision of the genus *Arcania* Leach, 1817 (Crustacea: Decapoda: Leucosioidea). *Zoologische Mededelingen, Leiden* 75: 169-206.
- Galil, B.S. 2001b. A revision of *Myra* Leach, 1817 (Crustacea: Decapoda: Leucosioidea). *Zoologische Mededelingen, Leiden* 75: 409-446.
- Garth, J.S. 1987. *Platypilumnus soelae*, a new species of goneplacid crab from the North West Shelf of Australia (Crustacea: Decapoda: Brachyura). *The Beagle, Occasional Papers of the Northern Territory Museum of Arts and Sciences* 4: 35-38.
- Garth, J.S., and Davie, P.J.F. 1995. A new species of *Parthenope* (Crustacea: Decapoda: Parthenopidae) from deep-water off northern Queensland. *Memoirs of the Queensland Museum* 38: 223-227.
- Goeke, G.D. 1985. Decapod Crustacea: Raninidae. In: *Résultats des Campagnes MUSORSTOM I et II - Philippines (1976, 1980)*. Tome 2. *Mémoires du Muséum National d'Histoire Naturelle, Paris A (Zoologie)* 133: 207-228.
- Grey, D.L., Dall, W., and Baker, A. 1983. *A guide to the Australian penaeid prawns*. Northern Territory Government Printing Office: Darwin. 140 pp.
- Griffin, D.J.G. 1970. The Australian majid spider crabs of the genus *Achaeus* (Crustacea, Brachyura). *Journal of the Royal Society of Western Australia* 53: 97-119.
- Griffin, D.J.G. 1972. Brachyura collected by Danish expeditions on south-eastern Australia (Crustacea, Decapoda). *Steenstrupia* 2: 49-90.
- Griffin, D.J.G. 1973. A revision of the spider crabs of the genus *Phalangipus* (Crustacea, Brachyura, Majidae). *Journal of Natural History* 7: 165-207.
- Griffin, D.J.G. 1976. Spider Crabs of the family Majidae (Crustacea: Brachyura) from the Philippine Islands. *Journal of Natural History* 10: 179-222.
- Griffin, D.J.G., and Tranter, H.A. 1986. The Decapoda Brachyura of the Siboga Expedition. Part VIII Majidae. *Siboga Expédition* 39: 1-335.
- Griffin, D.J.G., and Stoddart, H.E. 1995. Deep-water decapod Crustacea from eastern Australia: lobsters of the families Nephropidae, Palinuridae, Polychelidae and Scyllaridae. *Records of the Australian Museum* 47: 231-263.
- Guinot, D. 1989. Les genre *Carcinoplax* H. Milne Edwards, 1852 (Crustacea, Brachyura: Goneplacidae). In: Forest, J. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 5. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 144: 265-345.
- Guinot, D., and Richer de Forges, B. 1982. Révision du genre Indo-Pacifique *Cyrtomaia* Miers, 1886: Campagnes océanographiques du *Challenger*, de l'*Albatross*, du *Siboga* et du *Vauban* (Crustacea Decapoda Brachyura). *Annales de l'Institut Océanographique, Monaco et Paris* 58: 5-88.
- Guinot, D., and Richer de Forges, B. 1985. Crustacés Décapodes: Majidae (genres *Platymaia*, *Cyrtomaia*, *Pleistacantha*, *Sphenocarcinus* et *Naxioides*). In: *Résultats des Campagnes MUSORSTOM I et II - Philippines (1976, 1980)*. Tome 2. *Mémoires du Muséum National d'Histoire Naturelle, Paris A (Zoologie)* 133: 83-179.
- Guinot, D., and Richer de Forges, B. 1995. Crustacea Decapoda Brachyura: révision de la famille des Homolidae De Haan, 1839. In: Crosnier, A. (ed.), *Résultats des Campagnes MUSORSTOM*, vol. 13. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 163: 283-517.
- Haig, J. 1965. The Porcellanidae (Crustacea, Anomura) of Western Australia, with descriptions of four new Australian species. *Journal of the Royal Society of Western Australia* 48: 97:118.
- Haig, J. 1974. The anomuran crabs of Western Australia: their distribution in the Indian Ocean and adjacent seas. *Journal of the Marine Biological Association of India* 14: 443-451.
- Haig, J. 1981. Porcellanid crabs from the Indo-west Pacific, Part II. *Steenstrupia* 7: 269-291.
- Hale, H.M. 1927. *The crustaceans of South Australia. Part 1*. South Australian Government Printer: Adelaide. 201 pp.
- Hanamura, Y. 1987. Caridean shrimps obtained by R.V. "Soela" from north-west Australia, with description of a new species of *Leptochela* (Crustacea: Decapoda: Pasiphaeidae). *The*

- Beagle, Occasional Papers of the Northern Territory Museum of Arts and Sciences* 4: 15-33.
- Hanamura, Y., and Takeda, Y. 1987. Family Pandalidae (Crustacea, Decapoda, Caridea) collected by the RV "Soela" from the northwest Australian shelf. *Bulletin of the National Science Museum, Tokyo, Ser. A (Zoology)* 13: 103-121.
- Hanamura, Y., and Evans, D.R. 1994. Deepwater caridean shrimps of the families Oplophoridae and Pasiphaeidae (Crustacea: Decapoda) from Western Australia, with an appendix on a lophogastridan mysid (Mysidacea). *Crustacean Research* 23: 46-60.
- Hanamura, Y., and Evans, D.R. 1996. Deepwater caridean shrimps of the families Nematocarcinidae, Stylodactylidae, Pandalidae and Crangonidae (Crustacea: Decapoda) from Western Australia. *Bulletin of Nansei National Fisheries Research Institute* 29: 1-18.
- Hayashi, K.-I. 1975. The Indo-West Pacific Processidae (Crustacea, Decapoda, Caridea). *Journal of the Shimonoseki University of Fisheries* 24: 47-145.
- Hayashi, K.-I. 2004. Revision of the *Pasiphaea cristata* Bate, 1888 species group of *Pasiphaea* Savigny, 1816, with descriptions of four new species and referral of *P. australis* to *Alainopasiphaea* Hayashi, 1999 (Crustacea: Decapoda: Pasiphaeidae). In: Marshall, B.A., and Richer de Forges, B. (eds), Tropical Deep-Sea Benthos, Vol. 23. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 191: 319-373.
- Holthuis, L.B. 1946. Biological results of the Snellius Expedition XIV. The Decapoda Macrura of the Snellius Expedition I. The Stenopodidae, Nephropsidae, Scyllaridae and Palinuridae. *Temminckia* 7: 1-178, 11 pls.
- Holthuis, L.B. 1985. A revision of the family Scyllaridae (Crustacea: Decapoda: Macrura). I. Subfamily Ibaciniae. *Zoologische Verhandelingen, Leiden* 218: 1-130.
- Holthuis, L.B. 1991. FAO species catalogue. Vol. 13. Marine lobsters of the world. An annotated and illustrated catalogue of species of interest to fisheries known to date. *FAO Fisheries Synopsis* 125: 1-292.
- Holthuis, L.B. 1993. *The recent genera of the caridean and stenopodidean shrimps (Crustacea, Decapoda) with an appendix on the Order Amphionidacea*. Nationaal Natuurhistorisch Museum: Leiden. 328 pp.
- Holthuis, L.B. 2002. The Indo-Pacific scyllarine lobsters (Crustacea, Decapoda, Scyllaridae). *Zoosystema* 24: 499-683.
- Holthuis, L.B., and Manning, R.B. 1990. Crabs of the subfamily Dorippinae MacLeay, 1838, from the Indo-West Pacific region (Crustacea: Decapoda: Dorippidae). *Researches on Crustacea, Carcinological Society of Japan Special Number* 3: 1-151.
- Jones, D.S., and Morgan, G.J. 1993. An annotated checklist of the Crustacea of Rottnest Island, Western Australia. Pp. 135-162 in: Wells, F.E., Walker, D.I., Kirkman, H., and Lethbridge, R. (eds), *Proceedings of the Fifth International Marine Biological Workshop: The Marine Flora and Fauna of Rottnest Island, Western Australia*. Vol. 1. Western Australian Museum: Perth.
- Kensley, B. 1983. New records of bresiliid shrimp from Australia, South Africa, Caribbean, and Gulf of Mexico (Decapoda: Natantia: Caridea). *Smithsonian Contributions to Zoology* 394: 1-31.
- Kensley, B. 1989. New genera in the thalassinidean families Calocarididae and Axiidae (Crustacea: Decapoda). *Proceedings of the Biological Society of Washington* 102: 960-967.
- Kensley, B. 2003. Axioid shrimps from Guam (Crustacea, Decapoda, Thalassinidea). *Micronesica* 35-36: 359-384.
- Kensley, B., Tranter, H.A., and Griffin, D.J.G. 1987. Deepwater decapod Crustacea from eastern Australia (Penaeidea and Caridea). *Records of the Australian Museum* 39: 263-331.
- Koh, S.K., and Ng, P.K.L. 2000. A revision of the spiny crabs of the genus *Hypothalassia* Gistel, 1848 (Crustacea: Decapoda: Brachyura: Eriphiidae). *Raffles Bulletin Of Zoology* 48: 123-141.
- Komai, T. 2004. A review of the Indo-West Pacific species of the genus *Glyphocrangon* A. Milne-Edwards, 1881 (excluding the *G. caeca* species group) (Crustacea: Decapoda: Caridea: Glyphocrangonidae). In: Marshall, B.A., and Richer de Forges, B. (eds), Tropical Deep-Sea Benthos, Vol. 23. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 191: 375-610.
- Komai, T., and Chan, T.-Y. 2007. A new species of the crangonid shrimp genus *Philoceras* (Crustacea: Decapoda: Caridea) from the Philippines. *Proceedings of the Biological Society of Washington* 120: 159-166.
- Last, P.R., Lyne, V.D., Yearsley, G.K., Gledhill, D.C., Gomon, M.F., and Rees, A.J.J. 2005. *Validation of national demersal fish datasets for the regionalisation of the Australian continental slope and outer shelf (>40 m depth)*. The National Oceans Office and CSIRO Marine Research, Australia: Hobart. 98 pp.
- Leene, J.E. 1940. Biological results of the Snellius Expedition. VI. The Portunidae of the Snellius Expedition (Part 1). *Temminckia* 5: 163-188.
- Lemaitre, R. 1996. Hermit crabs of the family Parapaguridae (Crustacea: Decapoda: Anomura) from Australia: species of *Strobopagurus* Lemaitre, 1989, *Sympagurus* Smith, 1883 and two new genera. *Records of the Australian Museum* 48: 163-221.
- Lemaitre, R. 1999. Crustacea Decapoda: a review of the species of the genus *Parapagurus* Smith, 1879 (Parapaguridae) from the Pacific and Indian Oceans. In: Crosnier, A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 20. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 180: 303-378.
- Lemaitre, R. 2004a. A review of *Strobopagurus* Lemaitre, 1989 (Crustacea: Decapoda: Paguroidea: Parapaguridae), with description of a new species. *Scientia Marina* 68: 355-372.
- Lemaitre, R. 2004b. A worldwide review of hermit crab species of the genus *Sympagurus* Smith, 1883 (Crustacea: Decapoda: Parapaguridae). In: Marshall, B.A., and Richer de Forges, B. (eds), Tropical Deep-Sea Benthos, Vol. 23. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 191: 85-149.
- Lewinsohn, C. 1982. Eine neue Art der Gattung *Spiropagurus* Stimpson (Decapoda) aus dem Roten Meer. *Crustaceana* 42: 212-218.
- Lewis, M. 1999. CSIRO-SEBS (Seamount, Epibenthic Sampler), a new epibenthic sled for sampling seamounts and other rough terrain. *Deep Sea Research I. Oceanographic Research Papers* 46: 1101-1107.
- Lucas, J.S. 1980. Spider crabs of the family Hymenosomatidae (Crustacea; Brachyura) with particular reference to Australian species: systematics and biology. *Records of the Australian Museum* 33: 148-247.
- Macpherson, E. 1990. Crustacea Decapoda: on a collection of Nephropidae from the Indian Ocean and Western Pacific. In: Crosnier, A. (ed.), Résultats des Campagnes

- MUSORSTOM, vol. 6. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 145: 289-328.
- Macpherson, E. 1993a. Crustacea Decapoda: species of the genus *Paramunida* Baba, 1988 (Galatheidæ) from the Philippines, Indonesia and New Caledonia. In: Crosnier, A. (ed.), Résultats des Campagnes MUSORSTOM, vol. 10. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 156: 443-473.
- Macpherson, E. 1993b. New records for the genus *Nephropsis* Wood-Mason (Crustacea, Decapoda, Nephropidae) from northern Australia, with the description of two new species. *The Beagle, Occasional Papers of the Northern Territory Museum of Arts and Sciences* 10: 55-66.
- Macpherson, E. 1994. Crustacea Decapoda: Studies on the genus *Munida* Leach, 1820 (Galatheidæ) in New Caledonian and adjacent waters with descriptions of 56 new species. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 161: 421-569.
- Macpherson, E. 2007. Species of the genus *Munidopsis* Whiteaves, 1784 from the Indian and Pacific Oceans and reestablishment of the genus *Galacantha* A. Milne-Edwards, 1880 (Crustacea, Decapoda, Galatheidæ). *Zootaxa* 1417: 1-135.
- Macpherson, E., and Machordom, A. 2001. Phylogenetic relationships of species of *Raymunida* (Decapoda: Galatheidæ) based on morphology and mitochondrial cytochrome oxidase sequences, with the recognition of four new species. *Journal of Crustacean Biology* 21: 696-714.
- Man, J.G. de 1920. Pasiphaeidae, Stylodactylidae, Hoplophoridae, Nematocarcinidae, Thalassocaridae, Pandalidae, Psalidopodidae, Gnathophyllidae, Processidae, Glyphocranconidae and Crangonidae. *Siboga Expédition* 39: 1-318, pls 311-325.
- Manning, R.B., and Holthuis, L.B. 1981. West African brachyuran crabs. *Smithsonian Contributions to Zoology* 306: 1-379.
- McLaughlin, P.A. 1994. A new genus and two new species of deep-water hermit crabs (Decapoda: Anomura: Paguridae) from the Southern Ocean. *Proceedings of the Biological Society of Washington* 107: 469-481.
- McLaughlin, P.A. 1997. Crustacea Decapoda: hermit crabs of the family Paguridae from the KARUBAR cruise in Indonesia. In: Crosnier, A and Bouchet, P. (eds), Résultats des Campagnes MUSORSTOM, vol. 16. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 172: 433-572.
- McLaughlin, P.A. 2000. Crustacea Decapoda: *Porcellanopagurus* Filhol and *Solitariopagurus* Türkay (Paguridae), from New Caledonian area, Vanuatu and the Marquesas: new records, new species. In: Crosnier, A. (ed), Résultats des Campagnes MUSORSTOM, vol. 21. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 184: 389-414.
- McLaughlin, P.A. 2003. Illustrated keys to families and genera of the superfamily Paguroidea (Crustacea: Decapoda: Anomura), with diagnoses of genera of Paguridae. *Memoirs of Museum Victoria* 60: 111-144.
- McLaughlin, P.A. 2004. A review of the hermit crab genus *Nematopagurus* A. Milne-Edwards and Bouvier, 1892 (Crustacea: Decapoda: Paguridae). In: Marshall, B.A., and Richer de Forges, B. (eds), Tropical Deep-Sea Benthos, Vol. 23. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 191: 151-229.
- McLaughlin, P.A., and Haig, J. 1989. On the status of *Pylopaguropsis zebra* (Henderson), *P. magnimanus* (Henderson), and *Galapagurus teevanus* Boone, with descriptions of seven new species of *Pylopaguropsis* (Crustacea: Anomura: Paguridae). *Micronesica* 22: 123-171.
- McLaughlin, P.A., and Haig, J. 1996. A new genus for *Anapagrides* sensu de Saint Laurent-Dehancé, 1966 (Decapoda: Anomura: Paguridae) and descriptions of four new species. *Proceedings of the Biological Society of Washington* 109: 75-90.
- McLaughlin, P.A., and de Saint Laurent, M. 1998. A new genus for four species of hermit crabs formerly assigned to the genus *Pagurus* Fabricius (Decapoda: Anomura: Paguridae). *Proceedings of the Biological Society of Washington* 111: 158-187.
- McLay, C.L. 1993. Crustacea Decapoda: the sponge crabs (Dromiidae) of New Caledonia and the Philippines with a review of the genera. In: Crosnier, A. (ed.), Résultats de Campagnes MUSORSTOM, vol. 10. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 156: 111-251.
- McLay, C.L. 1999. Crustacea: Decapoda: Revision of the family Dynomenidae. In: Crosnier, A. (ed.), Résultats des Campagnes MUSORSTOM. Vol. 20. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 180: 427-569.
- Miers, E.J. 1876. On *Actaeomorpha erosa*, a new genus and species of Crustacea. *Journal of the Linnean Society of London (Zoology)* 13: 183-185, pl. 114.
- Miers, E.J. 1879. Crustacea. In: An account of the petrological, botanical, and zoological collections made in Kerguelen's Land and Rodriguez during the Transit of Venus Expeditions ... in the years 1874-75. *Philosophical Transactions of the Royal Society of London* 168: 200-214, pl. 211.
- Milne Edwards, H. 1837. *Histoire naturelle des crustacés, comprenant l'anatomie, la physiologie et la classification es ces animaux*. Vol. 2. Librairie Encyclopédique de Roret: Paris. 531 pp.
- Morgan, G.J. 1991. A review of the hermit crab genus *Calcinus* Dana (Crustacea: Decapoda: Diogenidae) from Australia, with descriptions of two new species. *Invertebrate Taxonomy* 5: 869-913.
- Morgan, G.J., and Forest, J. 1991. Seven new species of hermit crabs from Northern and Western Australia (Decapoda, Anomura, Diogenidae). *Bulletin du Muséum National d'Histoire Naturelle, Paris A*, 12: 649-689.
- Morgan, G.J., and Jones, D.S. 1991. Checklist of marine decapod Crustacea of southern Western Australia. Pp. 483-497 in: Wells, F.E., Walker, D.I., Kirkman, H., and Lethbridge, R. (eds), *Proceedings of the Third International Marine Biological Workshop: The Marine Flora and Fauna of Albany, Western Australia*. Vol. 2. Western Australian Museum: Perth.
- Ng, P.K.L. 1987. The Indo-Pacific Pilumnidae II. A revision of the genus *Rhizopa* Stimpson, 1858, and the status of the Rhizopinae Stimpson, 1858 (Crustacea, Decapoda, Brachyura). *Indo-Malayan Zoology* 4: 69-111.
- Ng, P.K.L. 1993. Krausiinae, a new subfamily for the genera *Kraussia* Dana, 1852, *Palpedia*, new genus, and *Garthasia* new genus (Crustacea: Decapoda: Brachyura: Xanthidae), with descriptions of two new species from Singapore and the Philippines. *Raffles Bulletin Of Zoology* 41: 133-157.
- Ng, P.K.L. 1996. *Garthambrus*, a new genus of deep water parthenopid crabs (Crustacea: Decapoda: Brachyura) from the Indo-Pacific, with description of a new species from the Seychelles. *Zoologische Mededelingen, Leiden* 70: 155-168.

- Ng, P.K.L. 1998. Crabs. Pp. 1045-1155 in: Carpenter, K.E., and Niem, V.H. (eds), *The living resources of the western central Pacific*. Vol. 2. Cephalopods, crustaceans, holothurians and sharks. FAO: Rome.
- Ng, P.K.L., and Chuang, C.T.N. 1996. The Hymenosomatidae (Crustacea: Decapoda: Brachyura) of southeast Asia, with notes on other species. *Raffles Bulletin of Zoology, Supplement 3*: 1-82.
- Ng, P.K.L., and Chan, T.-Y. 2000. Note on *Mathildella serrata* (Sakai, 1974) (Crustacea: Decapoda: Brachyura: Goneplacidae) from deep water in Taiwan. Pp. 149-153 in: Hwang, J.-S., Wang, C.-H., and Chan, T.-Y. (eds), *Proceedings of the International Symposium on Marine Biology in Taiwan -Crustacea and zooplankton taxonomy, ecology and living resources, 27-27 May, 1998, Taiwan*. National Taiwan Museum: Taipei.
- Ng, P.K.L., and Manuel-Santos, M.R. 2007. Establishment of the Vultocinidae, a new family for an unusual new genus and new species of Indo-West Pacific crab (Crustacea: Decapoda: Brachyura: Goneplacoidea), with comments on the taxonomy of the Goneplacidae. *Zootaxa* 1558: 39-68.
- Ng, P.K.L., Chan, T.-Y., and Wang, C.-H. 2000. The crabs of the families Dromiidae, Raninidae, Corystidae and Palicidae (Crustacea: Decapoda: Brachyura) of Taiwan. Pp. 155-180 in: Hwang, J.-S., Wang, C.-H., and Chan, T.-Y. (eds), *Proceedings of the International Symposium on Marine Biology in Taiwan -Crustacea and zooplankton taxonomy, ecology and living resources, 27-27 May, 1998, Taiwan*. National Taiwan Museum: Taipei.
- Ng, P.K.L., Guinot, D., and Davie, P.J.F. 2008. Systema Brachyurorum; Part I. An annotated checklist of extant brachyuran crabs of the world. *Raffles Bulletin Of Zoology* 17: 1-286.
- Ngoc-Ho, N. 1994. Some Callianassidae and Upogebidae from Australia with description of four new species (Crustacea: Decapoda: Thalassinidea). *Memoirs of the Museum of Victoria* 54: 51-78.
- Ngoc-Ho, N. 2006. Three species of *Acanthaxius* Sakai & de Saint Laurent, 1989, including two new to science, from the Solomon Islands and New Caledonia (Crustacea, Thalassinidea, Axiidae). *Zootaxa* 1240: 57-68.
- Noël, P. 1986. Crustacés Décapodes: Processidae de l'Indo-ouest-Pacifique. In: Forest, J. (ed.), *Résultats des Campagnes MUSORSTOM I & II, Philippines*, vol. 2. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 133: 261-301.
- O'Hara, T.D., and Poore, G.C.B. 2000. Patterns of distribution for southern Australian marine echinoderms and decapods. *Journal of Biogeography* 27: 1321-1335.
- Okuno, J. 1994. A new species of hinge-beak shrimp from the Western Pacific (Crustacea, Decapoda, Rhynchocinetidae). *The Beagle, Occasional Papers of the Northern Territory Museum of Arts and Sciences* 11: 29-37.
- Osawa, M., and Okuno, J. 2004. *Lauriea siagiani* Baba, 1994 (Crustacea, Decapoda, Anomura, Galatheidae) from Hachijo-jima Island, the Izu Islands, Japan. *Biogeography* 6: 57-62.
- Pérez Farfante, I. 1980. Revision of the penaeid shrimp genus *Penaeopsis* (Crustacea: Decapoda). *Fishery Bulletin of the Fish and Wildlife Service, United States* 77: 721-763.
- Pérez Farfante, I., and Kensley, B. 1997. Penaeoid and sergestoid shrimps and prawns of the world. Keys and diagnoses for the families and genera. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 175: 1-233.
- Poore, G.C.B. 1994. A phylogeny of the families of Thalassinidea (Crustacea: Decapoda) with keys to the families and genera. *Memoirs of the Museum of Victoria* 54: 79-120.
- Poore, G.C.B. 1997. A review of the thalassinidean families Callianideidae Kossman, Micheleidae Sakai, and Thomassiniidae de Saint Laurent (Crustacea: Decapoda) with descriptions of fifteen new species. *Zoosystema* 19: 345-420.
- Poore, G.C.B. 2004. *Marine decapod Crustacea of southern Australia. A guide to identification (with chapter on Stomatopoda by Shane Ahyong)*. CSIRO Publishing: Melbourne. 574 pp.
- Poore, G.C.B. in press. Thalassinidean shrimps (Crustacea: Decapoda) from north-western Australia, including five new species. *Records of the Western Australian Museum*.
- Poore, G.C.B., and Griffin, D.J.G. 1979. The Thalassinidea (Crustacea: Decapoda) of Australia. *Records of the Australian Museum* 32: 217-321.
- Poore, G.C.B., Hart, S., Taylor, J., and Tudge, C. 1998. Decapod crustaceans from Tasmanian seamounts. Pp. 65-78 in: Koslow, J.A., and Gowlett-Holmes, K. (eds), *The seamount fauna of southern Tasmania: benthic communities, their conservation and impacts of trawling. Final report to Environment Australia and The Fisheries Research Development Corporation*. CSIRO Marine Research: Hobart.
- Poupin, J., and McLaughlin, P.A. 1998. Additional records of *Calcinus* species (Decapoda: Anomura: Diogenidae) from French Polynesia with description of three new species and a key to Indo-West Pacific species of the genus. *Crustacean Research* 27: 9-27.
- Rahayu, D.L. 2006. The genus *Paguristes* (Crustacea, Decapoda, Diogenidae) from Indonesia. In: Richer de Forges, B., and Justine, J.-L. (eds), *Tropical Deep-Sea Benthos*, Vol. 24. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 193: 349-374.
- Rathbun, M.J. 1906. The Brachyura and Macrura of the Hawaiian Islands. *Bulletin of the United States Fisheries Commission* 23: 827-930. pls 821-824.
- Saint Laurent, M. de 1989. La nouvelle superfamille des Retroplumoidea Gill, 1894 (Decapoda, Brachyura): systématique, affinités et évolution. In: Forest, J. (ed.), *Résultats des Campagnes Musorstom*, vol. 5. *Mémoires du Muséum National d'Histoire Naturelle, Paris* 144: 103-179.
- Saint Laurent, M. de, and Cleve, R. 1981. Crustacés Décapodes : Stenopodidea. Pp. 151-188 in: *Résultats de Campagnes MUSORSTOM. 1. Philippines (18-28 mars 1976)*. Vol. 1. ORSTOM - Muséum National d'Histoire Naturelle: Paris.
- Sakai, K. 1976. *Crabs of Japan and adjacent seas*. Kodansha Ltd: Tokyo. 773 pp. plus volume of 251 plates pp.
- Sakai, K. 1982. Revision of Upogebiidae (Decapoda, Thalassinidea) in the Indo-West Pacific region. *Researches on Crustacea, Special Number* 1: 1-106.
- Sakai, K. 1986. *Axiopsis brucei* sp. nov., a new sponge-inhabiting axiid (Crustacea: Decapoda: Thalassinidea), from north-west Australia. *The Beagle, Occasional Papers of the Northern Territory Museum of Arts and Sciences* 3: 11-20.
- Sakai, K. 1988. A new genus and five new species of Callianassidae (Crustacea: Decapoda: Thalassinidea) from northern Australia. *The Beagle, Occasional Papers of the Northern Territory Museum of Arts and Sciences* 5: 51-69.
- Sakai, K. 1992. Axiid collections of the Zoological Museum, Copenhagen, with the description of one new genus and six

- new species (Axiidae, Thalassinidea, Crustacea). *Zoologica Scripta* 21: 157-180.
- Sakai, K. 1993. On a collection of Upogebiidae (Crustacea, Thalassinidea) from the Northern Territory Museum, Australia, with the description of two new species. *The Beagle, Occasional Papers of the Northern Territory Museum of Arts and Sciences* 10: 87-114.
- Sakai, K. 1994. Eleven species of Australian Axiidae (Crustacea: Decapoda: Thalassinidea) with descriptions of one new genus and five new species. *The Beagle, Occasional Papers of the Northern Territory Museum of Arts and Sciences* 11: 175-202.
- Sakai, K. 1999. Synopsis of the family Callianassidae, with keys to subfamilies, genera and species, and the description of new taxa (Crustacea: Decapoda: Thalassinidea). *Zoologische Verhandelingen, Leiden* 326: 1-152.
- Sakai, K. 2002. Callianassidae (Decapoda, Thalassinidea) in the Andaman Sea, Thailand. *Phuket Marine Biological Center Special Publication* 23: 461-532.
- Sakai, K. 2005. Callianassoidea of the world (Decapoda: Thalassinidea). *Crustaceana Monographs* 4: 1-285.
- Sakai, K., and Ohta, S. 2005. Some thalassinid collections by R/V "Hakuhou-Maru" and R/V "Tansei-Maru", University of Tokyo, in the Sulu Sea, Philippines, and in Sagami Bay and Suruga Bay, Japan, including two new species, one new genus, and one new family (Decapoda, Thalassinidea). *Crustaceana* 78: 67-93.
- Salva, E.W., and Feldmann, R.M. 2001. Reevaluation of the family Atelecyclidae (Decapoda: Brachyura). *Kirtlandia* 52: 9-62.
- Serène, R. 1984. Crustacés Décapodes Brachyours de l'Océan Indien Occidental et de la Mer Rouge, Xanthoidea: Xanthidae et Trapeziidae. Avec un addendum par Crosnier, A. : Carpiliidae et Menippidae. *Faune Tropicale* 24: 1-400.
- Serène, R., and Lohavanijaya, P. 1973. The Brachyura (Crustacea: Decapoda) collected by the Naga Expedition, including a review of the Homolidae. *Naga Report. Scientific Results of Marine Investigations of the South China Sea and the Gulf of Thailand* 4: 1-186, 186 figs, 121 pls.
- Short, J.W., and Davie, P.J.F. 1993. Decapoda Crustacea - non-Caridea. Pp. 67-69, Appendix 67 on p. 123 in: Russell, B.C., and Hanley, J.R. (eds), *The marine biological resources and heritage values of Cartier and Hibernia Reefs, Timor Sea*. Northern Territory Museum of Arts and Sciences: Darwin.
- Stephenson, W. 1972. An annotated check list and key to the Indo-West-Pacific swimming crabs (Crustacea: Decapoda: Portunidae). *Bulletin of the Royal Society of New Zealand* 10: 1-64.
- Tan, C.G.S. 1996. Leucosiidae of the *Albatross* expedition to the Philippines, 1907-1910 (Crustacea: Brachyura: Decapoda). *Journal of Natural History* 30: 1021-1058.
- Tan, C.G.S., and Ng, P.K.L. 1996. A revision of the Indo-Pacific genus *Oreophorus* Rüppell, 1830 (Crustacea: Decapoda: Brachyura: Leucosiidae). Pp. 101-189 in: Richer de Forges, B. (ed.) *Les fonds meubles des lagons de Nouvelle-Calédonie (sédimentologie, benthos)*. Vol. 2. ORSTOM Editions: Paris.
- Tan, S.H., Huang, J.F., and Ng, P.K.L. 2000. The deep-water crabs of the families Homolidae and Latreillidae (Crustacea: Decapoda: Brachyura) from Taiwan. Pp. 181-189 in: Hwang, J.-S., Wang, C.-H., and Chan, T.-Y. (eds), *Proceedings of the International Symposium on Marine Biology in Taiwan -Crustacea and zooplankton taxonomy, ecology and living resources, 27-27 May, 1998, Taiwan*. National Taiwan Museum: Taipei.
- Tavares, M. 1993. Crustacea Decapoda: Les Cyclodorippidae et Cymonomidae de l'Indo-Ouest Pacifique à l'exclusion du genre *Cymonomus*. In: Crosnier, A. (ed.), *Résultats des campagnes MUSORSTOM Vol. 10. Mémoires du Muséum National d'Histoire Naturelle, Paris* 156: 253-313.
- Tesch, J.J. 1918. Decapoda F. (Decapoda Brachyura continued) Goneplacidae and Pinnotheridae. *Siboga Expédition* 39c1: 149-296, pls 147-118.
- Tirmizi, N.M., and Javed, W. 1976. A new species of *Munida* from the Indian Ocean with a redescription of a syntype of *Munida spinulifera* Miers, 1884 (Decapoda, Galatheidea). *Crustaceana* 31: 81-89.
- Tyndale-Biscoe, M., and George, R.W. 1962. The Oxystomata and Gymnopleura (Crustacea, Brachyura) of Western Australia with descriptions of two new species from Western Australia and one from India. *Journal of the Royal Society of Western Australia* 45: 65-96.
- Vereshchaka, A.L. 2000. Revision of the genus *Sergia* (Decapoda: Dendrobranchiata: Sergestidae): taxonomy and distribution. *Galathea Reports* 18: 69-207, pls 202-205.
- Wadley, V., and Evans, D. 1992. Crustaceans from the Deepwater Trawl Fisheries of Western Australia. *Western Australian Museum*: 1-44.
- Wee, D.P.C., and Ng, P.K.L. 1995. Swimming crabs of the genera *Charybdis* De Haan, 1833, and *Thalamita* Latreille, 1829 (Crustacea: Decapoda: Brachyura: Portunidae) from Peninsular Malaysia and Singapore. *Raffles Bulletin of Zoology, Supplement* 1: 1-128.
- Williams, A.B. 1982. Revision of the genus *Latreillia* Roux (Brachyura: Homoloidea). *Quaderni del Laboratorio di Tecnologia della Pesca* 3: 227-255.

Index to families and genera

<i>Acanthaxius</i>	92	<i>Cosmonotus</i>	40
<i>Acanthephyra</i>	85	Crangonidae	81
<i>Achaeus</i>	57	<i>Crenarctus</i>	16
<i>Actaea</i>	74	<i>Cryptolutea</i>	65
<i>Actaeomorpha</i>	41	Cyclodorippidae	36
<i>Aegaeon</i>	81	Cymonomidae	36
Aethridae	41	<i>Cymonomus</i>	36
<i>Agononida</i>	18	<i>Cyrtomaia</i>	58
<i>Alainopasiphaea</i>	89	<i>Dagnaudus</i>	38
<i>Albunea</i>	23	<i>Dardanus</i>	24
Albuneidae	23	<i>Demania</i>	75
<i>Allogalathea</i>	18	<i>Diogenes</i>	25
Alpheidae	78	Diogenidae	24
<i>Alpheopsis</i>	78	<i>Discias</i>	80
<i>Alpheus</i>	78	<i>Dorhynchus</i>	58
<i>Ambiaxius</i>	94	<i>Dorippe</i>	44
<i>Anapagrides</i>	27	Dorippidae	44
<i>Anchistioides</i>	80	<i>Dorphinaxius</i>	93
Anchistioididae	80	<i>Drachiella</i>	41
<i>Arcania</i>	49	<i>Dromia</i>	37
Aristeidae	12	Dromiidae	37
<i>Aristeus</i>	12	<i>Dumea</i>	58
Atelecyclidae	43	Dynomenidae	38
<i>Atergatopsis</i>	74	<i>Ebalia</i>	50
<i>Aulacolambrus</i>	64	<i>Echinolatus</i>	67
<i>Austrodromidia</i>	37	<i>Engystenopus</i>	92
<i>Austrolibinia</i>	54	<i>Enriquea</i>	18
Axiidae	92	<i>Entomonyx</i>	60
<i>Axiopsis</i>	92	<i>Ephippias</i>	58
<i>Bathypaguropsis</i>	27	Epialtidae	54
<i>Bathypalaemonella</i>	80	<i>Epigodromia</i>	37
Bathypalaemonellidae	80	<i>Eplumula</i>	39
<i>Bathypilumnus</i>	65	<i>Ethusa</i>	45
Benthesicymidae	12	Ethusidae	45
<i>Benthesicymus</i>	12	<i>Ethusina</i>	45
<i>Bouvieraxius</i>	92	<i>Eualus</i>	84
Bresiliidae	80	Eugonatonotidae	82
<i>Caecopilumnus</i>	65	<i>Eugonatonotus</i>	82
<i>Calappa</i>	42	<i>Eumedonus</i>	65
Calappidae	42	<i>Eupasiphae</i>	89
<i>Calcinus</i>	24	Euryplacidae	46
<i>Callianassa</i>	94	<i>Fultodromia</i>	37
Callianassidae	94	<i>Galathea</i>	18
<i>Calocarides</i>	93	Galatheidae	18
Calocarididae	94	<i>Garthambrus</i>	64
<i>Calvactaea</i>	74	<i>Glyphocrangon</i>	83
<i>Camatopsis</i>	46	Glyphocrangonidae	83
<i>Camposcia</i>	58	<i>Gomeza</i>	44
Campylonotidae	80	Goneplacidae	47
<i>Campylonotus</i>	80	Gourretiidae	95
<i>Carcinoplax</i>	47	<i>Griffinia</i>	54
Carpiliidae	43	<i>Grypachaeus</i>	59
<i>Carpilius</i>	43	<i>Hadropenaeus</i>	15
<i>Cestopagurus</i>	27	<i>Halicarcinus</i>	57
<i>Charybdis</i>	67	<i>Haliporoides</i>	15
Chasmocarcinidae	46	<i>Haliporus</i>	15
Chirostylidae	17	<i>Hayashidonus</i>	89
<i>Chlorodiella</i>	74	<i>Hemipagurus</i>	27
<i>Chlorotocella</i>	87	<i>Heterocarpoides</i>	87
<i>Chlorotocus</i>	87	<i>Heterocarpus</i>	87
<i>Ciliopagurus</i>	24	<i>Heteropilumnus</i>	66
<i>Corallianassa</i>	94	<i>Heteroplax</i>	46
Corystidae	44	<i>Hexaplax</i>	47

Hexapodidae	47	<i>Myrine</i>	53
Hippolytidae	84	<i>Nanocassiope</i>	75
<i>Hirsutodynamene</i>	38	<i>Naxioides</i>	55
<i>Homoioplax</i>	73	<i>Nectocarcinus</i>	69
<i>Homola</i>	38	Nematocarcinidae	85
Homolidae	38	<i>Nematocarcinus</i>	85
<i>Homologenus</i>	38	<i>Nematopagurus</i>	28
<i>Hyastenus</i>	55	<i>Neodorippe</i>	44
<i>Hymenopenaeus</i>	15	<i>Neopalicus</i>	63
Hymenosomatidae	57	Nephropidae	34
<i>Hypothalassia</i>	45	<i>Nephropsis</i>	34
Hypothalassiidae	45	<i>Notonyx</i>	47
<i>Ibacus</i>	16	<i>Notosceles</i>	40
Inachidae	57	<i>Novactaea</i>	76
Iphiculidae	49	<i>Odontozona</i>	92
<i>Iphiculus</i>	49	<i>Oncinopus</i>	59
<i>Janicella</i>	85	<i>Oncopagurus</i>	31
<i>Jonas</i>	44	Oplophoridae	85
<i>Krangalangia</i>	36	<i>Oplophorus</i>	85
<i>Lahaina</i>	55	<i>Oreophorus</i>	53
<i>Latreillia</i>	39	<i>Ovalipes</i>	69
Latreilliidae	39	<i>Pachycheles</i>	22
<i>Latreillopsis</i>	39	Paguridae	27
<i>Lauriea</i>	19	<i>Paguristes</i>	25
<i>Lebbeus</i>	84	Palaemonidae	86
<i>Leptochela</i>	89	<i>Palapedia</i>	76
<i>Leptomithrax</i>	61	Palicidae	63
<i>Leucosia</i>	51	<i>Paliculus</i>	63
Leucosiidae	49	Palinuridae	16
<i>Libystes</i>	68	Pandalidae	87
<i>Liocarcinus</i>	68	Panopeidae	73
<i>Lipkecallianassa</i>	95	<i>Paractaea</i>	76
<i>Lissocarcinus</i>	68	<i>Paradorippe</i>	44
<i>Lissoporcellana</i>	22	<i>Paragiopagurus</i>	32
<i>Lithodes</i>	27	<i>Paralomis</i>	27
Lithodidae	27	<i>Paramolopsis</i>	39
<i>Lophopagurus</i>	28	<i>Paramunida</i>	21
<i>Lophoplax</i>	66	Parapaguridae	31
<i>Lupocyclus</i>	69	<i>Parapagurus</i>	32
<i>Lyreidus</i>	40	<i>Parapalicus</i>	63
<i>Lysmata</i>	84	<i>Parapenaeus</i>	13
<i>Maja</i>	61	<i>Parapontocaris</i>	81
Majidae	60	<i>Parapontophilus</i>	81
<i>Marianaxius</i>	93	<i>Paraselwynia</i>	66
<i>Mathildella</i>	48	<i>Parathranites</i>	69
Mathildellidae	47	<i>Paraxanthias</i>	77
<i>Medaeus</i>	75	<i>Paraxanthodes</i>	77
<i>Megaesthesius</i>	46	<i>Parilia</i>	53
<i>Merhippolyte</i>	84	<i>Parthenope</i>	64
<i>Merocryptus</i>	52	Parthenopidae	64
<i>Mertonia</i>	66	<i>Pasiphaea</i>	89
<i>Metacrangon</i>	81	Pasiphaeidae	89
<i>Metanephrops</i>	34	Penaeidae	13
<i>Metapenaeopsis</i>	13	<i>Penaeopsis</i>	13
<i>Michelea</i>	95	<i>Penaeus</i>	13
Micheleidae	95	<i>Pentacheles</i>	91
<i>Michelopagurus</i>	28	<i>Periclimenes</i>	86
<i>Micropalicus</i>	63	<i>Petrolisthes</i>	22
<i>Monodaeus</i>	75	<i>Phalangipus</i>	55
<i>Munida</i>	19	<i>Philocheras</i>	81
<i>Munidopsis</i>	21	<i>Philyra</i>	53
<i>Mursia</i>	42	<i>Phylladiorhynchus</i>	21
<i>Myra</i>	52	<i>Physachaeus</i>	59
Pilumnidae	65	<i>Pilumnopeus</i>	66

<i>Pilumnus</i>	66	Rhynchocinetidae	90
<i>Planotergum</i>	62	<i>Rochinia</i>	56
<i>Platylambrus</i>	65	<i>Sabinea</i>	82
<i>Platymaia</i>	60	Scyllaridae	16
<i>Platypilumnus</i>	47	<i>Sergestes</i>	14
<i>Platypodia</i>	77	Sergestidae	14
<i>Pleistacantha</i>	60	<i>Sergia</i>	14
<i>Plesionika</i>	87	<i>Sicyonia</i>	14
<i>Polycheles</i>	91	Sicyoniidae	14
Polychelidae	91	<i>Solenocera</i>	15
<i>Polyonyx</i>	23	Solenoceridae	15
<i>Pontocaris</i>	82	<i>Spiropagurus</i>	29
<i>Porcellanella</i>	23	<i>Stemonopa</i>	23
Porcellanidae	22	Stenopodidae	92
<i>Porcellanopagurus</i>	29	<i>Strigopagurus</i>	26
Portunidae	67	<i>Strobopagurus</i>	32
<i>Portunus</i>	70	<i>Sunipea</i>	60
<i>Prismatopus</i>	62	<i>Sympagurus</i>	33
<i>Processa</i>	89	<i>Synalpheus</i>	79
Processidae	89	<i>Systellaspis</i>	86
<i>Procletes</i>	88	<i>Takedromia</i>	37
<i>Propagurus</i>	29	<i>Tethisea</i>	95
<i>Pseudaristeus</i>	12	<i>Thalamita</i>	71
<i>Pseudolambrus</i>	65	Thalassocarididae	90
<i>Pseudopalicus</i>	64	<i>Thalassocaris</i>	90
<i>Psopheticus</i>	47	<i>Thyrolambrus</i>	65
<i>Puerulus</i>	16	<i>Tozeuma</i>	84
<i>Pycnoplax</i>	47	<i>Trachypenaeus</i>	13
<i>Pylocheles</i>	33	Trapeziidae	73
Pylochelidae	33	<i>Trichopeltarion</i>	43
<i>Pylopaguroopsis</i>	29	<i>Trigonoplax</i>	57
<i>Quadrella</i>	73	<i>Turleana</i>	30
<i>Randallia</i>	53	<i>Tymolus</i>	36
Raninidae	40	<i>Umalia</i>	41
<i>Raymunida</i>	21	<i>Upogebia</i>	96
<i>Remiarctus</i>	16	Upogebiidae	96
<i>Retropluma</i>	72	<i>Uroptychus</i>	17
Retroplumidae	72	Xanthidae	74
<i>Rhinolambrus</i>	65	<i>Yaldwynopsis</i>	39
<i>Rhynchocinetes</i>	90		



MUSEUMVICTORIA